

SNX1 Antibody (C-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP14044b

Specification

SNX1 Antibody (C-term) - Product Information

Application	WB,E
Primary Accession	O13596
Other Accession	O4R503 , NP_683758.1 , NP_690039.1 , NP_003090.2
Reactivity	Human
Predicted	Monkey
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	59070
Antigen Region	426-455

SNX1 Antibody (C-term) - Additional Information

Gene ID 6642

Other Names

Sorting nexin-1, SNX1

Target/Specificity

This SNX1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 426-455 amino acids from the C-terminal region of human SNX1.

Dilution

WB~~1:1000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

SNX1 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

SNX1 Antibody (C-term) - Protein Information

Name SNX1

Function Involved in several stages of intracellular trafficking. Interacts with membranes containing phosphatidylinositol 3-phosphate (PtdIns(3P)) or phosphatidylinositol 3,5-bisphosphate (PtdIns(3,5)P2) (PubMed:[12198132](#)). Acts in part as component of the retromer membrane-deforming SNX-BAR subcomplex. The SNX-BAR retromer mediates retrograde transport of cargo proteins from endosomes to the trans-Golgi network (TGN) and is involved in endosome-to-plasma membrane transport for cargo protein recycling. The SNX-BAR subcomplex functions to deform the donor membrane into a tubular profile called endosome-to-TGN transport carrier (ETC) (Probable). Can sense membrane curvature and has in vitro vesicle-to-membrane remodeling activity (PubMed:[19816406](#), PubMed:[23085988](#)). Involved in retrograde endosome-to-TGN transport of lysosomal enzyme receptors (IGF2R, M6PR and SORT1) and Shiginella dysenteria toxin stxB. Plays a role in targeting ligand-activated EGFR to the lysosomes for degradation after endocytosis from the cell surface and release from the Golgi (PubMed:[12198132](#), PubMed:[15498486](#), PubMed:[17101778](#), PubMed:[17550970](#), PubMed:[18088323](#), PubMed:[21040701](#)). Involvement in retromer-independent endocytic trafficking of P2RY1 and lysosomal degradation of protease-activated receptor-1/F2R (PubMed:[16407403](#), PubMed:[20070609](#)). Promotes KALRN- and RHO-dependent but retromer-independent membrane remodeling such as lamellipodium formation; the function is dependent on GEF activity of KALRN (PubMed:[20604901](#)). Required for endocytosis of DRD5 upon agonist stimulation but not for basal receptor trafficking (PubMed:[23152498](#)).

Cellular Location

Endosome membrane; Peripheral membrane protein; Cytoplasmic side. Golgi apparatus, trans-Golgi network membrane; Peripheral membrane protein; Cytoplasmic side. Early endosome membrane; Peripheral membrane protein; Cytoplasmic side. Cell projection, lamellipodium. Note=Enriched on tubular elements of the early endosome membrane. Binds preferentially to highly curved membranes enriched in phosphatidylinositol 3-phosphate (PtdIns(3P)) or phosphatidylinositol 3,5-bisphosphate (PtdIns(3,5)P2) (PubMed:[15498486](#)). Colocalized with SORT1 to tubular endosomal membrane structures called endosome-to-TGN transport carriers (ETCs) which are budding from early endosome vacuoles just before maturing into late endosome vacuoles (PubMed:[18088323](#)). Colocalizes with DNAJC13 and Shiginella dysenteria toxin stxB on early endosomes (PubMed:[19874558](#)) Colocalized with F-actin at the leading edge of lamellipodia in a KALRN-dependent manner (PubMed:[20604901](#)).


SNX1 Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

SNX1 Antibody (C-term) - Images





Western blot analysis of lysates from HeLa, 293T cell line (from left to right), using SNX1 Antibody (C-term)(Cat. #AP14044b). AP14044b was diluted at 1:1000 at each lane. A goat anti-rabbit IgG H&L(HRP) at 1:5000 dilution was used as the secondary antibody. Lysates at 35ug per lane.

SNX1 Antibody (C-term) - Background

This gene encodes a member of the sorting nexin family. Members of this family contain a phox (PX) domain, which is a phosphoinositide binding domain, and are involved in intracellular trafficking. This endosomal protein regulates the cell-surface expression of epidermal growth factor receptor. This protein also has a role in sorting protease-activated receptor-1 from early endosomes to lysosomes. This protein may form oligomeric complexes with family members. This gene results in three transcript variants encoding distinct isoforms.

SNX1 Antibody (C-term) - References

Nisar, S., et al. *Traffic* 11(4):508-519(2010)
Mari, M., et al. *Traffic* 9(3):380-393(2008)
Bryant, D.M., et al. *J. Cell. Sci.* 120 (PT 10), 1818-1828 (2007) :
Rojas, R., et al. *Mol. Cell. Biol.* 27(3):1112-1124(2007)
Nguyen, L.N., et al. *Clin. Cancer Res.* 12(23):6952-6959(2006)