

CD3Z Antibody (C-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP1413b

Specification

CD3Z Antibody (C-term) - Product Information

Application	WB, FC,E
Primary Accession	P20963
Other Accession	O9XSJ9
Reactivity	Human, Mouse
Predicted	Pig
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Antigen Region	113-141

CD3Z Antibody (C-term) - Additional Information

Gene ID 919

Other Names

T-cell surface glycoprotein CD3 zeta chain, T-cell receptor T3 zeta chain, CD247, CD247, CD3Z, T3Z, TCRZ

Target/Specificity

This CD3Z antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 113-141 amino acids from the C-terminal region of human CD3Z.

Dilution

WB~~1:4000-1:8000
FC~~1:25

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

CD3Z Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

CD3Z Antibody (C-term) - Protein Information

Name CD247

Synonyms CD3Z, T3Z, TCRZ

Function Part of the TCR-CD3 complex present on T-lymphocyte cell surface that plays an essential role in adaptive immune response. When antigen presenting cells (APCs) activate T-cell receptor (TCR), TCR- mediated signals are transmitted across the cell membrane by the CD3 chains CD3D, CD3E, CD3G and CD3Z. All CD3 chains contain immunoreceptor tyrosine-based activation motifs (ITAMs) in their cytoplasmic domain. Upon TCR engagement, these motifs become phosphorylated by Src family protein tyrosine kinases LCK and FYN, resulting in the activation of downstream signaling pathways (PubMed:[1384049](#), PubMed:[1385158](#), PubMed:[2470098](#), PubMed:[7509083](#)). CD3Z ITAMs phosphorylation creates multiple docking sites for the protein kinase ZAP70 leading to ZAP70 phosphorylation and its conversion into a catalytically active enzyme (PubMed:[7509083](#)). Plays an important role in intrathymic T-cell differentiation. Additionally, participates in the activity-dependent synapse formation of retinal ganglion cells (RGCs) in both the retina and dorsal lateral geniculate nucleus (dLGN) (By similarity).

Cellular Location

Cell membrane {ECO:0000250|UniProtKB:P24161}; Single-pass type I membrane protein

Tissue Location


CD3Z is expressed in normal lymphoid tissue and in peripheral blood mononuclear cells (PBMCs) (PubMed:11722641)

CD3Z Antibody (C-term) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


CD3Z Antibody (C-term) - Images



Western blot analysis of CD3Z (arrow) using rabbit polyclonal CD3Z Antibody (C-term) (Cat.#1413b).293 cell lysates (2 ug/lane) either nontransfected (Lane 1) or transiently transfected with the CD3Z gene (Lane 2) (Origene Technologies).



All lanes : Anti-CD3Z Antibody (C-term) at 1:4000-1:8000 dilution Lane 1: Jurkat whole cell lysate
Lane 2: MOLT-4 whole cell lysate Lane 3: Mouse thymus lysate Lysates/proteins at 20 µg per lane.
Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 19 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



Overlay histogram showing K562 cells stained with AP1413b(green line). The cells were fixed with 2% paraformaldehyde (10 min). The cells were then incubated in 2% bovine serum albumin to block non-specific protein-protein interactions followed by the antibody (AP1413b, 1:25 dilution) for 60 min at 37°C. The secondary antibody used was Goat-Anti-Rabbit IgG, DyLight® 488 Conjugated Highly Cross-Adsorbed(1583138) at 1/200 dilution for 40 min at 37°C. Isotype control antibody (blue line) was rabbit IgG1 (1µg/1x10⁶ cells) used under the same conditions. Acquisition of >10, 000 events was performed.

CD3Z Antibody (C-term) - Background

T-cell receptor zeta, together with T-cell receptor alpha/beta and gamma/delta heterodimers, and with CD3-gamma, -delta and -epsilon, forms the T-cell receptor-CD3 complex. The zeta chain plays an important role in coupling antigen recognition to several intracellular signal-transduction pathways. Low expression of the antigen results in impaired immune response.

CD3Z Antibody (C-term) - References

Miyagawa,H., Rheumatology (Oxford) 47 (2), 158-164 (2008)
Gorman,C.L., J. Immunol. 180 (2), 1060-1070 (2008)
Eleftheriadis,T., Am. J. Nephrol. 28 (1), 152-157 (2008)