

GOLGA5 Antibody (Center)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP14147c

Specification

GOLGA5 Antibody (Center) - Product Information

Application	WB, FC,E
Primary Accession	Q8TBA6
Other Accession	NP_005104.2
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	83024
Antigen Region	381-408

GOLGA5 Antibody (Center) - Additional Information

Gene ID 9950

Other Names

Golgin subfamily A member 5, Cell proliferation-inducing gene 31 protein, Golgin-84, Protein Ret-II, RET-fused gene 5 protein, GOLGA5, RETII, RFG5

Target/Specificity

This GOLGA5 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 381-408 amino acids from the Central region of human GOLGA5.

Dilution

WB~~1:1000
FC~~1:10~50

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

GOLGA5 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

GOLGA5 Antibody (Center) - Protein Information

Name GOLGA5

Synonyms RETII, RFG5

Function Involved in maintaining Golgi structure. Stimulates the formation of Golgi stacks and ribbons. Involved in intra-Golgi retrograde transport.

Cellular Location

Golgi apparatus membrane; Single-pass type IV membrane protein. Note=Found throughout the Golgi, both on cisternae and, at higher abundance, on the tubulo-vesicular structures of the cis-Golgi network

Tissue Location


Ubiquitous. Highly expressed in seminiferous tubules and Leydig cells in testis, and detected at much lower levels in the other tissues tested. Expression is very low or not detectable in spermatozoa.

GOLGA5 Antibody (Center) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

GOLGA5 Antibody (Center) - Images



GOLGA5 Antibody (Center) (Cat. #AP14147c) western blot analysis in K562 cell line lysates (35ug/lane). This demonstrates the GOLGA5 antibody detected the GOLGA5 protein (arrow).



GOLGA5 Antibody (Center) (Cat. #AP14147c) flow cytometric analysis of K562 cells (right histogram) compared to a negative control cell (left histogram). FITC-conjugated donkey-anti-rabbit secondary antibodies were used for the analysis.

GOLGA5 Antibody (Center) - Background

The Golgi apparatus, which participates in glycosylation and transport of proteins and lipids in the secretory pathway, consists of a series of stacked cisternae (flattened membrane sacs). Interactions between the Golgi and microtubules are thought to be important for the reorganization of the Golgi after it fragments during mitosis. This gene encodes one of the golgins, a family of proteins localized to the Golgi. This protein is a coiled-coil membrane protein that has been postulated to play a role in vesicle tethering and docking. Translocations involving this gene and the ret proto-oncogene have been found in tumor tissues; the chimeric sequences have been designated RET-II and PTC5.

GOLGA5 Antibody (Center) - References

- Olsen, J.V., et al. Cell 127(3):635-648(2006)
- Olsen, J.V., et al. Cell 127(3):635-648(2006)
- Malsam, J., et al. Science 307(5712):1095-1098(2005)
- Rush, J., et al. Nat. Biotechnol. 23(1):94-101(2005)
- Brill, L.M., et al. Anal. Chem. 76(10):2763-2772(2004)