

FZD5 Antibody (C-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP14210b

Specification

FZD5 Antibody (C-term) - Product Information

Application	WB,E
Primary Accession	Q13467
Other Accession	NP_003459.2
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	64507
Antigen Region	472-500

FZD5 Antibody (C-term) - Additional Information

Gene ID 7855

Other Names

Frizzled-5, Fz-5, hFz5, FzE5, FZD5, C2orf31

Target/Specificity

This FZD5 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 472-500 amino acids from the C-terminal region of human FZD5.

Dilution

WB~~1:1000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

FZD5 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

FZD5 Antibody (C-term) - Protein Information

Name FZD5

Synonyms C2orf31

Function Receptor for Wnt proteins (PubMed:[10097073](#), PubMed:[20530549](#), PubMed:[26908622](#), PubMed:[9054360](#)). Functions in the canonical Wnt/beta- catenin signaling pathway. In vitro activates WNT2, WNT10B, WNT5A, but not WNT2B or WNT4 signaling (By similarity). In neurons, activation by WNT7A promotes formation of synapses (PubMed:[20530549](#)). May be involved in transduction and intercellular transmission of polarity information during tissue morphogenesis and/or in differentiated tissues (Probable). Plays a role in yolk sac angiogenesis and in placental vascularization (By similarity). Plays a role in ocular development (PubMed:[26908622](#)).

Cellular Location


Cell membrane; Multi-pass membrane protein {ECO:0000250|UniProtKB:Q8CHL0}. Golgi apparatus membrane {ECO:0000250|UniProtKB:Q9EQD0}; Multi-pass membrane protein {ECO:0000250|UniProtKB:Q9EQD0}. Synapse {ECO:0000250|UniProtKB:Q8CHL0}. Perikaryon {ECO:0000250|UniProtKB:Q8CHL0}. Cell projection, dendrite {ECO:0000250|UniProtKB:Q8CHL0}. Cell projection, axon {ECO:0000250|UniProtKB:Q8CHL0}. Note=Localized at the plasma membrane and also found at the Golgi apparatus. {ECO:0000250|UniProtKB:Q9EQD0}

FZD5 Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

FZD5 Antibody (C-term) - Images



FZD5 Antibody (C-term) (Cat. #AP14210b) western blot analysis in A549 cell line lysates (35ug/lane). This demonstrates the FZD5 antibody detected the FZD5 protein (arrow).

FZD5 Antibody (C-term) - Background

Members of the 'frizzled' gene family encode 7-transmembrane domain proteins that are receptors for Wnt signaling proteins. The FZD5 protein is believed to be the receptor for the Wnt5A ligand.

FZD5 Antibody (C-term) - References

- Moller, M., et al. BMC Infect. Dis. 10, 154 (2010) :
Yerges, L.M., et al. J. Bone Miner. Res. 24(12):2039-2049(2009)
Terabayashi, T., et al. J. Biol. Chem. 284(39):26716-26724(2009)
Kim, J.G., et al. J. Korean Med. Sci. 24(3):443-447(2009)
Lin, Z., et al. J. Biol. Chem. 283(48):33053-33058(2008)