

ALDH2 Antibody
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP1432d

Specification

ALDH2 Antibody - Product Information

Application	WB, IHC-P, FC,E
Primary Accession	P05091
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Antigen Region	1-517

ALDH2 Antibody - Additional Information

Gene ID 217

Other Names

Aldehyde dehydrogenase, mitochondrial, ALDH class 2, ALDH-E2, ALDH1, ALDH2, ALDM

Target/Specificity

This ALDH2 antibody is generated from rabbits immunized with human ALDH2 recombinant protein.

Dilution

WB~~1:1000
IHC-P~~1:50~100
FC~~1:10~50

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

ALDH2 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

ALDH2 Antibody - Protein Information

Name ALDH2

Synonyms ALDM

Function Required for clearance of cellular formaldehyde, a cytotoxic and carcinogenic

metabolite that induces DNA damage.

Cellular Location


Mitochondrion matrix.

ALDH2 Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

ALDH2 Antibody - Images




ALDH2 Antibody (Cat. #AP1432d) western blot analysis in A549,HepG2 cell line lysates (35ug/lane).This demonstrates the ALDH2 antibody detected the ALDH2 protein (arrow).



Formalin-fixed and paraffin-embedded human brain tissue reacted with ALDH2 Antibody, which

was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.



ALDH2 Antibody (Cat. #AP1432d) flow cytometry analysis of MDA-MB231 cells (bottom histogram) compared to a negative control cell (top histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

ALDH2 Antibody - Background

ALDH2 belongs to the aldehyde dehydrogenase family of proteins. Aldehyde dehydrogenase is the second enzyme of the major oxidative pathway of alcohol metabolism. Two major liver isoforms of this enzyme, cytosolic and mitochondrial, can be distinguished by their electrophoretic mobilities, kinetic properties, and subcellular localizations. Most Caucasians have two major isozymes, while approximately 50% of Asians have only the cytosolic isozyme, missing the mitochondrial isozyme. A remarkably higher frequency of acute alcohol intoxication among Asians than among Caucasians could be related to the absence of the mitochondrial isozyme.

ALDH2 Antibody - References

- Guo, Y.M., World J. Gastroenterol. 14 (9), 1444-1449 (2008)
- Chen, L., PLoS Med. 5 (3), E52 (2008)
- Teeguarden, J.G., Inhal Toxicol 20 (4), 375-390 (2008)
- Yoshida, A., Pharmacogenetics 2 (4), 139-147 (1992)