

**TRIP10 Antibody (N-term)**  
**Affinity Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP14358A**

**Specification**

---

**TRIP10 Antibody (N-term) - Product Information**

Application	WB,E
Primary Accession	<a href="#">Q15642</a>
Other Accession	<a href="#">P97531</a> , <a href="#">Q8CJ53</a> , <a href="#">NP_004231.1</a>
Reactivity	Human
Predicted	Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	68352
Antigen Region	115-144

**TRIP10 Antibody (N-term) - Additional Information**

**Gene ID** 9322

**Other Names**

Cdc42-interacting protein 4, Protein Felic, Salt tolerant protein, hSTP, Thyroid receptor-interacting protein 10, TR-interacting protein 10, TRIP-10, TRIP10, CIP4, STOT, STP

**Target/Specificity**

This TRIP10 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 115-144 amino acids from the N-terminal region of human TRIP10.

**Dilution**

WB~~1:1000

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

TRIP10 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

**TRIP10 Antibody (N-term) - Protein Information**

**Name** TRIP10

**Synonyms** CIP4, STOT, STP

**Function** Required for translocation of GLUT4 to the plasma membrane in response to insulin signaling (By similarity). Required to coordinate membrane tubulation with reorganization of the actin cytoskeleton during endocytosis. Binds to lipids such as phosphatidylinositol 4,5-bisphosphate and phosphatidylserine and promotes membrane invagination and the formation of tubules. Also promotes CDC42-induced actin polymerization by recruiting WASL/N-WASP which in turn activates the Arp2/3 complex. Actin polymerization may promote the fission of membrane tubules to form endocytic vesicles. Required for the formation of podosomes, actin-rich adhesion structures specific to monocyte-derived cells. May be required for the lysosomal retention of FASLG/FASL.

**Cellular Location**

Cytoplasm, cytoskeleton. Cytoplasm, cell cortex. Lysosome. Golgi apparatus. Cell membrane. Cell projection, phagocytic cup. Note=Translocates to the plasma membrane in response to insulin stimulation, and this may require active RHOQ (By similarity) Localizes to cortical regions coincident with F-actin, to lysosomes and to sites of phagocytosis in macrophages. Also localizes to the Golgi, and this requires AKAP9.


**Tissue Location**

Expressed in brain, colon, heart, kidney, liver, lung, megakaryocyte, ovary, pancreas, peripheral blood lymphocytes, placenta, prostate, skeletal muscle, small intestine, spleen, testis, thymus and trachea.

**TRIP10 Antibody (N-term) - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**TRIP10 Antibody (N-term) - Images**

TRIP10 Antibody (N-term) (Cat. #AP14358a) western blot analysis in K562 cell line lysates

(35ug/lane). This demonstrates the TRIP10 antibody detected the TRIP10 protein (arrow).

#### **TRIP10 Antibody (N-term) - Background**

Required for translocation of GLUT4 to the plasma membrane in response to insulin signaling (By similarity). Required to coordinate membrane tubulation with reorganization of the actin cytoskeleton during endocytosis. Binds to lipids such as phosphatidylinositol 4,5-bisphosphate and phosphatidylserine and promotes membrane invagination and the formation of tubules. Also promotes CDC42-induced actin polymerization by recruiting WASL/N-WASP which in turn activates the Arp2/3 complex. Actin polymerization may promote the fission of membrane tubules to form endocytic vesicles. Required for the formation of podosomes, actin-rich adhesion structures specific to monocyte-derived cells. May be required for the lysosomal retention of FASLG/FASL.

#### **TRIP10 Antibody (N-term) - References**

Roignot, J., et al. *Cancer Lett.* 288(1):116-123(2010)  
Hu, J., et al. *Cell. Signal.* 21(11):1686-1697(2009)  
Banerjee, P.P., et al. *J. Exp. Med.* 204(10):2305-2320(2007)  
Sugiyama, N., et al. *Mol. Cell Proteomics* 6(6):1103-1109(2007)  
Shimada, A., et al. *Cell* 129(4):761-772(2007)