

Mouse Acvr1c Antibody (C-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP14606b**Specification**

Mouse Acvr1c Antibody (C-term) - Product Information

Application	WB,E
Primary Accession	Q8K348
Other Accession	P70539 , Q8NER5 , NP_001104500.1
Reactivity	Mouse
Predicted	Human, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	54700
Antigen Region	341-368

Mouse Acvr1c Antibody (C-term) - Additional Information**Gene ID** 269275**Other Names**

Activin receptor type-1C, Activin receptor type IC, ACTR-IC, Activin receptor-like kinase 7, ALK-7, Acvr1c {ECO:0000312|EMBL:AAH287801}

Target/Specificity

This Mouse Acvr1c antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 341-368 amino acids from the C-terminal region of mouse Acvr1c.

Dilution

WB~~1:1000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Mouse Acvr1c Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Mouse Acvr1c Antibody (C-term) - Protein Information**Name** Acvr1c {ECO:0000312|EMBL:AAH28780.1}

Function Serine/threonine protein kinase which forms a receptor complex on ligand binding. The receptor complex consisting of 2 type II and 2 type I transmembrane serine/threonine kinases. Type II receptors phosphorylate and activate type I receptors which autophosphorylate, then bind and activate SMAD transcriptional regulators, SMAD2 and SMAD3. Receptor for activin AB, activin B and NODAL. Plays a role in cell differentiation, growth arrest and apoptosis.

Cellular Location

Membrane; Single-pass type I membrane protein

Tissue Location


Expressed in interdigital regions in developing limb buds.

Mouse Acvr1c Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Mouse Acvr1c Antibody (C-term) - Images



Mouse Acvr1c Antibody (C-term) (Cat. #AP14606b) western blot analysis in mouse heart tissue lysates (35ug/lane). This demonstrates the Acvr1c antibody detected the Acvr1c protein (arrow).

Mouse Acvr1c Antibody (C-term) - Background

Serine/threonine protein kinase which forms a receptor complex on ligand binding. The receptor complex consisting of 2 type II and 2 type I transmembrane serine/threonine kinases. Type II receptors phosphorylate and activate type I receptors which autophosphorylate, then bind and activate SMAD transcriptional regulators, SMAD2 and SMAD3. Receptor for activin AB, activin B and NODAL. Plays a role in cell differentiation, growth arrest and apoptosis.

Mouse Acvr1c Antibody (C-term) - References

Andersson, O., et al. Proc. Natl. Acad. Sci. U.S.A. 105(20):7252-7256(2008)

Bertolino, P., et al. Proc. Natl. Acad. Sci. U.S.A. 105(20):7246-7251(2008)

Liguori, G.L., et al. Dev. Biol. 315(2):280-289(2008)

Kurrasch, D.M., et al. J. Neurosci. 27(50):13624-13634(2007)

Kogame, M., et al. J. Med. Invest. 53 (3-4), 238-245 (2006) :

Mouse Acvr1c Antibody (C-term) - Citations

- [ALK7 protects against pathological cardiac hypertrophy in mice.](#)