

PLD2 Antibody (N-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP14669a

Specification

PLD2 Antibody (N-term) - Product Information

Application	WB,E
Primary Accession	O14939
Other Accession	Q0V8L6 , NP_002654.3
Reactivity	Human, Mouse
Predicted	Bovine
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Antigen Region	10-39

PLD2 Antibody (N-term) - Additional Information

Gene ID 5338

Other Names

Phospholipase D2, PLD 2, hPLD2, Choline phosphatase 2, PLD1C, Phosphatidylcholine-hydrolyzing phospholipase D2, PLD2

Target/Specificity

This PLD2 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 10-39 amino acids from the N-terminal region of human PLD2.

Dilution

WB~~1:2000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

PLD2 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

PLD2 Antibody (N-term) - Protein Information

Name PLD2 ([HGNC:9068](#))

Function Function as phospholipase selective for phosphatidylcholine (PubMed:[9582313](#)). May

have a role in signal-induced cytoskeletal regulation and/or endocytosis (By similarity).

Cellular Location

Cell membrane {ECO:0000250|UniProtKB:P97813}; Lipid-anchor {ECO:0000250|UniProtKB:P97813}


Tissue Location

Ubiquitous..


PLD2 Antibody (N-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

PLD2 Antibody (N-term) - Images


PLD2 Antibody (N-term) (Cat. #AP14669a) western blot analysis in mouse bladder tissue lysates (35ug/lane). This demonstrates the PLD2 antibody detected the PLD2 protein (arrow).



PLD2 Antibody (N-term) (Cat. #AP14669a) western blot analysis in MDA-MB453 cell line lysates (35ug/lane). This demonstrates the PLD2 antibody detected the PLD2 protein (arrow).



Western blot analysis of lysate from human heart tissue lysate, using PLD2 Antibody (N-term)(Cat. #AP14669a). AP14669a was diluted at 1:1000. A goat anti-rabbit IgG H&L(HRP) at 1:5000 dilution was used as the secondary antibody. Lysate at 35ug.



Anti-PLD2 Antibody (N-term) at 1:2000 dilution + Mouse lung tissue lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 106 kDa Blocking/Dilution buffer: 5% NFD/MTBST.

PLD2 Antibody (N-term) - Background

Phosphatidylcholine (PC)-specific phospholipases D (PLDs; EC 3.1.4.4) catalyze the hydrolysis of PC to produce phosphatidic acid and choline. Activation of PC-specific PLDs occurs as a consequence of agonist stimulation of both tyrosine kinase and G protein-coupled receptors. PC-specific PLDs have been proposed to function in regulated secretion, cytoskeletal reorganization, transcriptional regulation, and cell cycle control.[supplied by OMIM].

PLD2 Antibody (N-term) - References

Chae, Y.C., et al. Mol. Cell. Biol. 30(21):5086-5098(2010)
Bailey, S.D., et al. Diabetes Care 33(10):2250-2253(2010)
Knapek, K., et al. Mol. Cell. Biol. 30(18):4492-4506(2010)
Tabatabaian, F., et al. J. Biol. Chem. 285(25):18991-19001(2010)
Kang, D.W., et al. PLoS ONE 5 (8), E12109 (2010) :

PLD2 Antibody (N-term) - Citations

- [D-series Resolvins activate Phospholipase D in phagocytes during inflammation and resolution](#)
- [Phospholipase D1 Ablation Disrupts Mouse Longitudinal Hippocampal Axis Organization and Functioning](#)
- [Oxidized LDL phagocytosis during foam cell formation in atherosclerotic plaques relies on a PLD2-CD36 functional interdependence.](#)
- [AQP3 small interfering RNA and PLD2 small interfering RNA inhibit the proliferation and promote the apoptosis of squamous cell carcinoma.](#)