

COBRA1 Antibody (Center)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP14765c

Specification

COBRA1 Antibody (Center) - Product Information

Application	WB,E
Primary Accession	O8WX92
Other Accession	O8C4Y3 , NP_056271.2
Reactivity	Human
Predicted	Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	65697
Antigen Region	256-285

COBRA1 Antibody (Center) - Additional Information

Gene ID 25920

Other Names

Negative elongation factor B, NELF-B, Cofactor of BRCA1, NELFB, COBRA1, KIAA1182

Target/Specificity

This COBRA1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 256-285 amino acids from the Central region of human COBRA1.

Dilution

WB~~1:1000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

COBRA1 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

COBRA1 Antibody (Center) - Protein Information

Name NELFB

Synonyms COBRA1 {ECO:0000303|PubMed:11739404}, KI

Function Essential component of the NELF complex, a complex that negatively regulates the elongation of transcription by RNA polymerase II (PubMed:[12612062](#)). The NELF complex, which acts via an association with the DSIF complex and causes transcriptional pausing, is counteracted by the P-TEFb kinase complex (PubMed:[10199401](#)). May be able to induce chromatin unfolding (PubMed:[11739404](#)). Essential for early embryogenesis; plays an important role in maintaining the undifferentiated state of embryonic stem cells (ESCs) by preventing unscheduled expression of developmental genes (By similarity). Plays a key role in establishing the responsiveness of stem cells to developmental cues; facilitates plasticity and cell fate commitment in ESCs by establishing the appropriate expression level of signaling molecules (By similarity). Supports the transcription of genes involved in energy metabolism in cardiomyocytes; facilitates the association of transcription initiation factors with the promoters of the metabolism- related genes (By similarity).

Cellular Location

Nucleus.

Tissue Location


Widely expressed. Expressed in heart, brain, lung, placenta, liver, skeletal muscle, kidney and pancreas

COBRA1 Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

COBRA1 Antibody (Center) - Images



COBRA1 Antibody (Center) (Cat. #AP14765c) western blot analysis in 293 cell line lysates (35ug/lane). This demonstrates the COBRA1 antibody detected the COBRA1 protein (arrow).

COBRA1 Antibody (Center) - Background

NELFB is a subunit of negative elongation factor (NELF),

which also includes NELFA (WHSC2; MIM 606026), either NELFC or NELFD (TH1L; MIM 605297), and NELFE (RDBP; MIM 154040). NELF acts with DRB sensitivity-inducing factor (DSIF), a heterodimer of SPT4 (SUPT4H1; MIM 603555) and SPT5 (SUPT5H; MIM 602102), to cause transcriptional pausing of RNA polymerase II (see MIM 180660) (Narita et al., 2003 [PubMed 12612062]).

COBRA1 Antibody (Center) - References

Sun, J., et al. J. Cell. Biochem. 103(6):1798-1807(2008)
Sun, J., et al. J. Steroid Biochem. Mol. Biol. 107 (3-5), 131-139 (2007) :
Wu, C., et al. Proteomics 7(11):1775-1785(2007)
Narita, T., et al. Mol. Cell 26(3):349-365(2007)
Aiyar, S.E., et al. Int. J. Biol. Sci. 3(7):486-492(2007)