

NANOG Antibody (S285)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP1486E

Specification

NANOG Antibody (S285) - Product Information

Application	IF, WB,E
Primary Accession	O9H9S0
Other Accession	O6NSW7 , NP_079141
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	34620
Antigen Region	267-292

NANOG Antibody (S285) - Additional Information

Gene ID 79923

Other Names

Homeobox protein NANOG, Homeobox transcription factor Nanog, hNanog, NANOG

Target/Specificity

This NANOG antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 267-292 amino acids from human NANOG.

Dilution

IF~~1:10~50

WB~~1:1000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

NANOG Antibody (S285) is for research use only and not for use in diagnostic or therapeutic procedures.

NANOG Antibody (S285) - Protein Information

Name NANOG

Function Transcription regulator involved in inner cell mass and embryonic stem (ES) cells

proliferation and self-renewal. Imposes pluripotency on ES cells and prevents their differentiation towards extraembryonic endoderm and trophectoderm lineages. Blocks bone morphogenetic protein-induced mesoderm differentiation of ES cells by physically interacting with SMAD1 and interfering with the recruitment of coactivators to the active SMAD transcriptional complexes. Acts as a transcriptional activator or repressor. Binds optimally to the DNA consensus sequence 5'-TAAT[GT][GT]-3' or 5'-[CG][GA][CG]C[GC]ATTAN[GC]- 3'. Binds to the POU5F1/OCT4 promoter (PubMed:25825768). Able to autorepress its expression in differentiating (ES) cells: binds to its own promoter following interaction with ZNF281/ZFP281, leading to recruitment of the NuRD complex and subsequent repression of expression. When overexpressed, promotes cells to enter into S phase and proliferation.

Cellular Location

Nucleus {ECO:0000255|PROSITE-ProRule:PRU00108, ECO:0000269|PubMed:15983365}

Tissue Location


Expressed in testicular carcinoma and derived germ cell tumors (at protein level). Expressed in fetal gonads, ovary and testis. Also expressed in ovary teratocarcinoma cell line and testicular embryonic carcinoma. Not expressed in many somatic organs and oocytes.

NANOG Antibody (S285) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

NANOG Antibody (S285) - Images



Fluorescent confocal image of SY5Y cells stained with AP1486e NANOG (S285) antibody. SY5Y cells were fixed with 4% PFA (20 min), permeabilized with Triton X-100 (0.2%, 30 min), then incubated with AP1486e NANOG (S285) primary antibody (1:200, 2 h at room temperature). For

secondary antibody, Alexa Fluor® 488 conjugated donkey anti-rabbit antibody (green) was used (1:1000, 1h). Cytoplasmic actin was counterstained with Alexa Fluor® 555 (red) conjugated Phalloidin (5.25 µM, 25 min). Nuclei were counterstained with Hoechst 33342 (blue) (10 µg/ml, 3 min). Nanog immunoreactivity is localized mainly to the nuclei and also to the cytoplasm.



All lanes : Anti-NANOG Antibody (S285) at 1:1000 dilution Lane 1: MCF-7 whole cell lysate Lane 2: human testis lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 35 kDa Blocking/Dilution buffer: 5% NFD/MTBST.

NANOG Antibody (S285) - Background

NANOG is a Transcription regulator involved in inner cell mass and embryonic stem (ES) cells proliferation and self-renewal. It imposes pluripotency on ES cells and prevents their differentiation towards extraembryonic endoderm and trophectoderm lineages. This protein blocks bone morphogenetic protein-induced mesoderm differentiation of ES cells by physically interacting with SMAD1 and interfering with the recruitment of coactivators to the active SMAD transcriptional complexes. NANOG acts as a transcriptional activator or repressor. It binds optimally to the DNA consensus sequence 5'-[CG][GA][CG]C[GC]ATTAN[GC]-3'. When overexpressed, this protein promotes cells to enter into S phase and proliferation.

NANOG Antibody (S285) - References

Kochupurakkal, B.S., *Biochem. Biophys. Res. Commun.* 365 (4), 846-850 (2008)
Freberg, C.T., *Mol. Biol. Cell* 18 (5), 1543-1553 (2007)