

ICOSLG Antibody (C-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP14872b

Specification

ICOSLG Antibody (C-term) - Product Information

Application	WB,E
Primary Accession	O75144
Other Accession	NP_056074.1
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	33349
Antigen Region	217-246

ICOSLG Antibody (C-term) - Additional Information

Gene ID 23308

Other Names

ICOS ligand, B7 homolog 2, B7-H2, B7-like protein GI50, B7-related protein 1, B7RP-1, CD275, ICOSLG, B7H2, B7RP1, ICOSL, KIAA0653

Target/Specificity

This ICOSLG antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 217-246 amino acids from the C-terminal region of human ICOSLG.

Dilution

WB~~1:1000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

ICOSLG Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

ICOSLG Antibody (C-term) - Protein Information

Name ICOSLG

Synonyms B7H2, B7RP1, ICOSL, KIAA0653

Function Ligand for the T-cell-specific cell surface receptor ICOS. Acts as a costimulatory signal for T-cell proliferation and cytokine secretion; induces also B-cell proliferation and differentiation into plasma cells. Could play an important role in mediating local tissue responses to inflammatory conditions, as well as in modulating the secondary immune response by co-stimulating memory T-cell function (By similarity).

Cellular Location

Cell membrane; Single-pass type I membrane protein

Tissue Location


Isoform 1 is widely expressed (brain, heart, kidney, liver, lung, pancreas, placenta, skeletal muscle, bone marrow, colon, ovary, prostate, testis, lymph nodes, leukocytes, spleen, thymus and tonsil), while isoform 2 is detected only in lymph nodes, leukocytes and spleen. Expressed on activated monocytes and dendritic cells.

ICOSLG Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

ICOSLG Antibody (C-term) - Images



ICOSLG Antibody (C-term) (Cat. #AP14872b) western blot analysis in HL-60 cell line lysates (35ug/lane). This demonstrates the ICOSLG antibody detected the ICOSLG protein (arrow).

ICOSLG Antibody (C-term) - Background

Ligand for the T-cell-specific cell surface receptor ICOS. Acts as a costimulatory signal for T-cell proliferation and cytokine secretion; induces also B-cell proliferation and differentiation into plasma cells. Could play an important role in mediating local tissue responses to inflammatory conditions, as well as in modulating the secondary immune response by co-stimulating memory T-cell function (By similarity).

ICOSLG Antibody (C-term) - References

- Shimada, M., et al. Hum. Genet. 128(4):433-441(2010)
Bailey, S.D., et al. Diabetes Care 33(10):2250-2253(2010)
Wang, K., et al. Hum. Mol. Genet. 19(10):2059-2067(2010)
Pan, F., et al. Immunogenetics 62(4):237-251(2010)
Dubois, P.C., et al. Nat. Genet. 42(4):295-302(2010)