

DKK2 Antibody (C-term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP1522b**Specification**

DKK2 Antibody (C-term) - Product Information

Application	WB, IHC-P,E
Primary Accession	O9UBU2
Other Accession	O9OYZ8
Reactivity	Human
Predicted	Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	28447
Antigen Region	225-251

DKK2 Antibody (C-term) - Additional Information**Gene ID** 27123**Other Names**

Dickkopf-related protein 2, Dickkopf-2, Dkk-2, hDkk-2, DKK2

Target/Specificity

This DKK2 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 225-251 amino acids from the C-terminal region of human DKK2.

DilutionWB~~1:1000
IHC-P~~1:50~100**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

DKK2 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

DKK2 Antibody (C-term) - Protein Information**Name** DKK2

Function Antagonizes canonical Wnt signaling by inhibiting LRP5/6 interaction with Wnt and by forming a ternary complex with the transmembrane protein KREMEN that promotes internalization of LRP5/6. DKKs play an important role in vertebrate development, where they locally inhibit Wnt regulated processes such as antero-posterior axial patterning, limb development, somitogenesis and eye formation. In the adult, Dkks are implicated in bone formation and bone disease, cancer and Alzheimer disease (By similarity).

Cellular Location

Secreted.

Tissue Location


Expressed in heart, brain, skeletal muscle and lung

DKK2 Antibody (C-term) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

DKK2 Antibody (C-term) - Images



DKK2 Antibody (C239) (Cat. #AP1522b) western blot analysis in 293 cell line lysates (35ug/lane). This demonstrates the DKK2 antibody detected the DKK2 protein (arrow).



Formalin-fixed and paraffin-embedded human cancer tissue reacted with the primary antibody, which was peroxidase-conjugated to the secondary antibody, followed by AEC staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated. BC = breast carcinoma; HC = hepatocarcinoma.

DKK2 Antibody (C-term) - Background

The 259-amino acid DKK2 protein, like DKK1, DKK3, and DKK4, possesses an N-terminal signal peptide and 2 conserved cysteine-rich domains, which are separated by a linker region and contain 10 cys residues each. The second cys region has a putative lipid-binding function that may facilitate WNT/DKK interactions at the plasma membrane. The linker region contains 50 to 55 amino acids in DKK1, DKK2, and DKK4, whereas in DKK3 it contains only 12 amino acids. All DKKs have several potential sites for cleavage by furin-type proteases. Northern blot analysis revealed expression of 4.0- and 4.5-kb DKK2 transcripts in heart, brain, skeletal muscle, and lung. Western blot analysis showed that DKK2 is secreted as a 15- to 17-kD protein. Functional analysis determined that DKK2 does not block *Xenopus* Wnt8 induction of a secondary axis in frog embryos.

DKK2 Antibody (C-term) - References

Clark, H.F., et al., *Genome Res.* 13(10):2265-2270 (2003).
Brott, B.K., et al., *Mol. Cell. Biol.* 22(17):6100-6110 (2002).
Krupnik, V.E., et al., *Gene* 238(2):301-313 (1999).