

DKK4 Antibody (C-term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP1524b

Specification

DKK4 Antibody (C-term) - Product Information

Application	WB, IHC-P,E
Primary Accession	O9UBT3
Other Accession	NP_055235
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	24876
Antigen Region	188-217

DKK4 Antibody (C-term) - Additional Information

Gene ID 27121

Other Names

Dickkopf-related protein 4, Dickkopf-4, Dkk-4, hDkk-4, Dickkopf-related protein 4 short form, DKK4

Target/Specificity

This DKK4 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 188-217 amino acids from the C-terminal region of human DKK4.

Dilution

WB~~1:1000
IHC-P~~1:50~100

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

DKK4 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

DKK4 Antibody (C-term) - Protein Information

Name DKK4

Function Antagonizes canonical Wnt signaling by inhibiting LRP5/6 interaction with Wnt and by

forming a ternary complex with the transmembrane protein KREMEN that promotes internalization of LRP5/6. DKKs play an important role in vertebrate development, where they locally inhibit Wnt regulated processes such as antero-posterior axial patterning, limb development, somitogenesis and eye formation. In the adult, Dkks are implicated in bone formation and bone disease, cancer and Alzheimer disease (By similarity).

Cellular Location

Secreted.

Tissue Location


Expressed in cerebellum, T-cells, esophagus and lung

DKK4 Antibody (C-term) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

DKK4 Antibody (C-term) - Images



The anti-DKK4 C-term Pab (Cat. #AP1524b) is used in Western blot to detect DKK4 in A375 cell lysate.



Formalin-fixed and paraffin-embedded human cancer tissue reacted with the primary antibody, which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated. BC = breast carcinoma; HC = hepatocarcinoma.

DKK4 Antibody (C-term) - Background

DKK4, like DKK1, DKK2, and DKK3, possesses an N-terminal signal peptide and 2 conserved cysteine-rich domains, which are separated by a linker region and contain 10 cysteine residues each. The second cysteine region has a putative lipid-binding function that may facilitate WNT/DKK interactions at the plasma membrane. The linker region contains 50 to 55 amino acids in DKK1, DKK2, and DKK4, whereas in DKK3 it contains only 12 amino acids. All DKKs have several potential sites for cleavage by furin-type proteases. Northern blot analysis detected no expression of DKK4, but RT-PCR analysis detected DKK4 expression in cerebellum, T-cell, esophagus, and lung cDNA libraries. DKK4 blocks *Xenopus* Wnt8, Wnt3a, and Wnt2b, but not Dsh or Fz8, induction of a secondary axis in frog embryos, indicating that DKKs antagonize WNT function upstream of WNT receptors.

DKK4 Antibody (C-term) - References

Krupnik, V.E., et al., *Gene* 238(2):301-313 (1999).
Yoshida, S., et al., *Cytogenet. Cell Genet.* 94 (1-2), 88-89 (2001).

DKK4 Antibody (C-term) - Citations

- [Expression pattern of REIC/Dkk-3 in various cell types and the implications of the soluble form in prostatic acinar development.](#)