

NFIC Antibody (Center)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP16052c

Specification

NFIC Antibody (Center) - Product Information

| | |
|-------------------|---|
| Application | IF, WB,E |
| Primary Accession | P08651 |
| Other Accession | NP_995315.1 , NP_005588.2 |
| Reactivity | Human |
| Host | Rabbit |
| Clonality | Polyclonal |
| Isotype | Rabbit IgG |
| Calculated MW | 55675 |
| Antigen Region | 180-208 |

NFIC Antibody (Center) - Additional Information

Gene ID 4782

Other Names

Nuclear factor 1 C-type, NF1-C, Nuclear factor 1/C, CCAAT-box-binding transcription factor, CTF, Nuclear factor I/C, NF-I/C, NFI-C, TGGCA-binding protein, NFIC, NFI

Target/Specificity

This NFIC antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 180-208 amino acids from the Central region of human NFIC.

Dilution

IF~~1:10~50
WB~~1:1000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

NFIC Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

NFIC Antibody (Center) - Protein Information

Name NFIC

Synonyms NFI

Function Recognizes and binds the palindromic sequence 5'- TTGGCNNNNNGCCAA-3' present in viral and cellular promoters and in the origin of replication of adenovirus type 2. These proteins are individually capable of activating transcription and replication.

Cellular Location


Nucleus.

NFIC Antibody (Center) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

NFIC Antibody (Center) - Images



Fluorescent confocal image of HeLa cell stained with NFIC Antibody (Center)(Cat#AP16052c). HeLa cells were fixed with 4% PFA (20 min), permeabilized with Triton X-100 (0.1%, 10 min), then incubated with NFIC primary antibody (1:25, 1 h at 37°C). For secondary antibody, Alexa Fluor® 488 conjugated donkey anti-rabbit antibody (green) was used (1:400, 50 min at 37°C). Cytoplasmic actin was counterstained with Alexa Fluor® 555 (red) conjugated Phalloidin (7units/ml, 1 h at 37°C). Nuclei were counterstained with DAPI (blue) (10 µg/ml, 10 min). NFIC immunoreactivity is localized to nucleus significantly.



All lanes : Anti-NFIC Antibody (Center) at 1:1000 dilution Lane 1: HeLa whole cell lysates Lane 2: A549 whole cell lysates Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution Predicted band size : 56 kDa Blocking/Dilution buffer: 5% NFDN/TBST.

NFIC Antibody (Center) - Background

NFIC recognizes and binds the palindromic sequence 5'-TTGGCNNNNNGCCAA-3' present in viral and cellular promoters and in the origin of replication of adenovirus type 2. These proteins are individually capable of activating transcription and replication.

NFIC Antibody (Center) - References

Rose, J.E., et al. Mol. Med. 16 (7-8), 247-253 (2010) :
Nilsson, J., et al. Cancer Res. 70(5):2020-2029(2010)
Udelhoven, M., et al. J. Mol. Endocrinol. 44(2):99-113(2010)
Plasari, G., et al. Mol. Cell. Biol. 29(22):6006-6017(2009)
Riffel, A.K., et al. Mol. Pharmacol. 76(5):1104-1114(2009)