

**UBA6 Antibody (C-term)**  
**Affinity Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP16886b****Specification**

---

**UBA6 Antibody (C-term) - Product Information**

Application	WB,E
Primary Accession	<a href="#">A0AVT1</a>
Other Accession	<a href="#">Q8C7R4</a> , <a href="#">NP_060697.4</a>
Reactivity	Human
Predicted	Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	117970
Antigen Region	861-889

**UBA6 Antibody (C-term) - Additional Information**

**Gene ID** 55236

**Other Names**

Ubiquitin-like modifier-activating enzyme 6, Ubiquitin-activating enzyme 6, Monocyte protein 4, MOP-4, Ubiquitin-activating enzyme E1-like protein 2, E1-L2, UBA6, MOP4, UBE1L2

**Target/Specificity**

This UBA6 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 861-889 amino acids from the C-terminal region of human UBA6.

**Dilution**

WB~~1:1000

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

UBA6 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

**UBA6 Antibody (C-term) - Protein Information**

**Name** UBA6

**Synonyms** MOP4, UBE1L2

**Function** Activates ubiquitin by first adenylating its C-terminal glycine residue with ATP, and thereafter linking this residue to the side chain of a cysteine residue in E1, yielding a ubiquitin-E1 thioester and free AMP. Specific for ubiquitin, does not activate ubiquitin-like peptides. Differs from UBE1 in its specificity for substrate E2 charging. Does not charge cell cycle E2s, such as CDC34. Essential for embryonic development. Required for UBD/FAT10 conjugation. Isoform 2 may play a key role in ubiquitin system and may influence spermatogenesis and male fertility.


**Tissue Location**

Widely expressed. Isoform 2 is predominantly expressed in testis with higher expression in adult testis than in fetal testis.

**UBA6 Antibody (C-term) - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**UBA6 Antibody (C-term) - Images**

UBA6 Antibody (C-term) (Cat. #AP16886b) western blot analysis in K562 cell line lysates (35ug/lane). This demonstrates the UBA6 antibody detected the UBA6 protein (arrow).

**UBA6 Antibody (C-term) - Background**

Modification of proteins with ubiquitin (UBB; MIM 191339) or ubiquitin-like proteins controls many signaling networks and requires a ubiquitin-activating enzyme (E1), a ubiquitin conjugating enzyme (E2), and a ubiquitin protein ligase (E3). UBE1L2 is an E1 enzyme that initiates the activation and conjugation of ubiquitin-like proteins (Jin et al., 2007 [PubMed 17597759]).

**UBA6 Antibody (C-term) - References**

Groettrup, M., et al. Trends Biochem. Sci. 33(5):230-237(2008)

Chiu, Y.H., et al. Mol. Cell 27(6):1014-1023(2007)

Pelzer, C., et al. J. Biol. Chem. 282(32):23010-23014(2007)

Jin, J., et al. Nature 447(7148):1135-1138(2007)

Hillier, L.W., et al. Nature 434(7034):724-731(2005)

**UBA6 Antibody (C-term) - Citations**

- [The non-canonical ubiquitin activating enzyme UBA6 suppresses epithelial-mesenchymal transition of mammary epithelial cells.](#)