

NOX1 Antibody (Center)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP17191c**Specification**

NOX1 Antibody (Center) - Product Information

Application	WB, IHC-P-Leica,E
Primary Accession	O9Y5S8
Other Accession	NP_008983.2 , NP_039249.1
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Antigen Region	243-271

NOX1 Antibody (Center) - Additional Information**Gene ID** 27035**Other Names**

NADPH oxidase 1, NOX-1, 1---, Mitogenic oxidase 1, MOX-1, NADH/NADPH mitogenic oxidase subunit P65-MOX, NOH-1, NOX1, MOX1, NOH1

Target/Specificity

This NOX1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 243-271 amino acids from the Central region of human NOX1.

DilutionWB~~1:1000
IHC-P-Leica~~1:500**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

NOX1 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

NOX1 Antibody (Center) - Protein Information**Name** NOX1 ([HGNC:7889](#))**Synonyms** MOX1, NOH1

Function NADPH oxidase that catalyzes the generation of superoxide from molecular oxygen utilizing NADPH as an electron donor.

Cellular Location

Cell projection, invadopodium membrane; Multi-pass membrane protein. Cell membrane; Multi-pass membrane protein

Tissue Location


[Isoform NOH-1L]: Detected in colon, uterus, prostate, and colon carcinoma, but not in peripheral blood leukocytes

NOX1 Antibody (Center) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


NOX1 Antibody (Center) - Images



All lanes : Anti-NOX1 Antibody (Center) at 1:1000 dilution Lane 1: WiDr whole cell lysate Lane 2: SW480 whole cell lysate Lane 3: HT-29 whole cell lysate Lane 4: PC-3 whole cell lysate Lane 5: HeLa whole cell lysate Lane 6: human uterus tissue lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 65 kDa Blocking/Dilution buffer: 5% NFD/MTBST.



Immunohistochemical analysis of paraffin-embedded human colon tissue using AP17191C performed on the Leica® BOND RXm. Tissue was fixed with formaldehyde at room temperature, antigen retrieval was by heat mediation with a EDTA buffer (pH9. 0). Samples were incubated with primary antibody(1:500) for 1 hours at room temperature. A undiluted biotinylated CRF Anti-Polyvalent HRP Polymer antibody was used as the secondary antibody.



Immunohistochemical analysis of paraffin-embedded human kidney tissue using AP17191C performed on the Leica® BOND RXm. Tissue was fixed with formaldehyde at room temperature, antigen retrieval was by heat mediation with a EDTA buffer (pH9. 0). Samples were incubated with primary antibody(1:500) for 1 hours at room temperature. A undiluted biotinylated CRF Anti-Polyvalent HRP Polymer antibody was used as the secondary antibody.

NOX1 Antibody (Center) - Background

Voltage-gated proton (hydrogen) channels play an important role in cellular defense against acidic stress. They are unique among ion channels with respect to their extremely high selectivity, marked temperature dependence, and unitary conductance, which is 3 orders of magnitude lower than that of most other ion channels. NOX1 is a homolog of the catalytic subunit of the superoxide-generating NADPH oxidase of phagocytes, gp91phox. Two transcript variants encoding different isoforms have been found

for this gene.

NOX1 Antibody (Center) - References

- Sancho, P., et al. J. Biol. Chem. 285(32):24815-24824(2010)
van Bruggen, R., et al. Blood 115(26):5398-5400(2010)
Roberts, K.E., et al. Gastroenterology 139(1):130-139(2010)
Malec, V., et al. Free Radic. Biol. Med. 48(12):1626-1635(2010)
Manea, A., et al. Biochem. Biophys. Res. Commun. 396(4):901-907(2010)