

GBP5 Antibody (N-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP17262a**Specification**

GBP5 Antibody (N-term) - Product Information

Application	WB,E
Primary Accession	O96PP8
Other Accession	NP_443174.1 , NP_001127958.1
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	66617
Antigen Region	140-167

GBP5 Antibody (N-term) - Additional Information**Gene ID** 115362**Other Names**

Guanylate-binding protein 5, GBP-TA antigen, GTP-binding protein 5, GBP-5, Guanine nucleotide-binding protein 5, GBP5

Target/Specificity

This GBP5 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 140-167 amino acids from the N-terminal region of human GBP5.

Dilution

WB~~1:1000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

GBP5 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

GBP5 Antibody (N-term) - Protein Information**Name** GBP5 {ECO:0000303|PubMed:20180847, ECO:0000312|HGNC:HGNC:19895}**Function** Interferon (IFN)-inducible GTPase that plays important roles in innate immunity against

a diverse range of bacterial, viral and protozoan pathogens (By similarity). Hydrolyzes GTP, but in contrast to other family members, does not produce GMP (PubMed:[20180847](#)). Following infection, recruited to the pathogen-containing vacuoles or vacuole- escaped bacteria and acts as a positive regulator of inflammasome assembly by promoting the release of inflammasome ligands from bacteria (By similarity). Acts by promoting lysis of pathogen-containing vacuoles, releasing pathogens into the cytosol (By similarity). Following pathogen release in the cytosol, promotes recruitment of proteins that mediate bacterial cytolysis: this liberates ligands that are detected by inflammasomes, such as lipopolysaccharide (LPS) that activates the non-canonical CASP4/CASP11 inflammasome or double- stranded DNA (dsDNA) that activates the AIM2 inflammasome (By similarity). As an activator of NLRP3 inflammasome assembly: promotes selective NLRP3 inflammasome assembly in response to microbial and soluble, but not crystalline, agents (PubMed:[22461501](#)). Independently of its GTPase activity, acts as an inhibitor of various viruses infectivity, such as HIV-1, Zika and influenza A viruses, by inhibiting FURIN-mediated maturation of viral envelope proteins (PubMed:[26996307](#), PubMed:[31091448](#)).

Cellular Location

Cytoplasmic vesicle membrane {ECO:0000250|UniProtKB:Q8CFB4}; Lipid-anchor, GPI-like-anchor. Golgi apparatus membrane; Lipid-anchor. Cytoplasm

Tissue Location


Expressed in peripheral blood monocytes (at protein level).

GBP5 Antibody (N-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

GBP5 Antibody (N-term) - Images



GBP5 Antibody (N-term) (Cat. #AP17262a) western blot analysis in T47D cell line lysates (35ug/lane). This demonstrates the GBP5 antibody detected the GBP5 protein (arrow).

GBP5 Antibody (N-term) - Background

Belongs to the GBP family.

GBP5 Antibody (N-term) - References

- Davila, S., et al. Genes Immun. 11(3):232-238(2010)
Wehner, M., et al. FEBS J. 277(7):1597-1605(2010)
Ito, Y., et al. J. Infect. Dis. 197(5):663-666(2008)
Tripal, P., et al. J. Interferon Cytokine Res. 27(1):44-52(2007)
Fellenberg, F., et al. J. Invest. Dermatol. 122(6):1510-1517(2004)