

MUM1 Antibody (C-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP17700b

Specification

MUM1 Antibody (C-term) - Product Information

Application	WB,E
Primary Accession	Q2TAK8
Other Accession	NP_116242.2
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	78636
Antigen Region	671-699

MUM1 Antibody (C-term) - Additional Information

Gene ID 84939

Other Names

PWWP domain-containing protein MUM1, Mutated melanoma-associated antigen 1, MUM-1, Protein expandere, MUM1, EXPAND1

Target/Specificity

This MUM1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 671-699 amino acids from the C-terminal region of human MUM1.

Dilution

WB~~1:1000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

MUM1 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

MUM1 Antibody (C-term) - Protein Information

Name PWWP3A ([HGNC:29641](#))

Function Involved in the DNA damage response pathway by contributing to the maintenance of

chromatin architecture. Recruited to the vicinity of DNA breaks by TP53BP1 and plays an accessory role to facilitate damage-induced chromatin changes and promoting chromatin relaxation. Required for efficient DNA repair and cell survival following DNA damage.

Cellular Location


Nucleus. Note=Recruited to DNA damage sites via its interaction with the BRCT domain of TP53BP1

MUM1 Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

MUM1 Antibody (C-term) - Images



MUM1 Antibody (C-term) (Cat. #AP17700b) western blot analysis in CEM cell line lysates (35ug/lane). This demonstrates the MUM1 antibody detected the MUM1 protein (arrow).

MUM1 Antibody (C-term) - Background

Involved in the DNA damage response pathway by contributing to the maintenance of chromatin architecture. Recruited to the vicinity of DNA breaks by TP53BP1 and plays an accessory role to facilitate damage-induced chromatin changes and promoting chromatin relaxation. Required for efficient DNA repair and cell survival following DNA damage.

MUM1 Antibody (C-term) - References

- Huen, M.S., et al. Mol. Cell 37(6):854-864(2010)
Matsuoka, S., et al. Science 316(5828):1160-1166(2007)
Coulie, P.G., et al. Proc. Natl. Acad. Sci. U.S.A. 92(17):7976-7980(1995)