

**ATG4A Antibody**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP1808c**

**Specification**

---

**ATG4A Antibody - Product Information**

Application	<b>WB, IHC-P,E</b>
Primary Accession	<a href="#">O8WYN0</a>
Other Accession	<a href="#">O6PZ05</a>
Reactivity	<b>Human, Mouse</b>
Predicted	<b>Bovine</b>
Host	<b>Rabbit</b>
Clonality	<b>Polyclonal</b>
Isotype	<b>Rabbit IgG</b>
Antigen Region	<b>363-392</b>

**ATG4A Antibody - Additional Information**

**Gene ID** 115201

**Other Names**

Cysteine protease ATG4A, 3422-, AUT-like 2 cysteine endopeptidase, Autophagin-2, Autophagy-related cysteine endopeptidase 2, Autophagy-related protein 4 homolog A, hAPG4A, ATG4A, APG4A, AUTL2

**Target/Specificity**

This ATG4A antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 363-392 amino acids from human ATG4A.

**Dilution**

WB~~1:1000  
IHC-P~~1:50~100

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

ATG4A Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

**ATG4A Antibody - Protein Information**

**Name** ATG4A {ECO:0000303|Ref.20, ECO:0000312|HGNC:HGNC:16489}

**Function** Cysteine protease that plays a key role in autophagy by mediating both proteolytic activation and delipidation of ATG8 family proteins (PubMed:[12473658](#), PubMed:[15169837](#), PubMed:[17347651](#), PubMed:[21177865](#), PubMed:[21245471](#), PubMed:[22302004](#), PubMed:[32732290](#)). The protease activity is required for proteolytic activation of ATG8 family proteins: cleaves the C-terminal amino acid of ATG8 proteins to reveal a C-terminal glycine (PubMed:[12473658](#), PubMed:[15169837](#), PubMed:[17347651](#), PubMed:[21177865](#), PubMed:[21245471](#), PubMed:[22302004](#)). Exposure of the glycine at the C-terminus is essential for ATG8 proteins conjugation to phosphatidylethanolamine (PE) and insertion to membranes, which is necessary for autophagy (PubMed:[12473658](#), PubMed:[15169837](#), PubMed:[17347651](#), PubMed:[21177865](#), PubMed:[21245471](#), PubMed:[22302004](#)). Preferred substrate is GABARAPL2 followed by MAP1LC3A and GABARAP (PubMed:[12473658](#), PubMed:[15169837](#), PubMed:[17347651](#), PubMed:[21177865](#), PubMed:[21245471](#), PubMed:[22302004](#)). Protease activity is also required to counteract formation of high-molecular weight conjugates of ATG8 proteins (ATG8ylation): acts as a deubiquitinating- like enzyme that removes ATG8 conjugated to other proteins, such as ATG3 (PubMed:[31315929](#), PubMed:[33773106](#)). In addition to the protease activity, also mediates delipidation of ATG8 family proteins (PubMed:[29458288](#), PubMed:[33909989](#)). Catalyzes delipidation of PE- conjugated forms of ATG8 proteins during macroautophagy (PubMed:[29458288](#), PubMed:[33909989](#)). Compared to ATG4B, the major protein for proteolytic activation of ATG8 proteins, shows weaker ability to cleave the C-terminal amino acid of ATG8 proteins, while it displays stronger delipidation activity (PubMed:[29458288](#)). Involved in phagophore growth during mitophagy independently of its protease activity and of ATG8 proteins: acts by regulating ATG9A trafficking to mitochondria and promoting phagophore-endoplasmic reticulum contacts during the lipid transfer phase of mitophagy (PubMed:[33773106](#)).

#### **Cellular Location**

Cytoplasm {ECO:0000250|UniProtKB:Q8BGE6}.


#### **ATG4A Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### **ATG4A Antibody - Images**






Western blot analysis of anti-APG4A Pab (Cat. #AP1808c) in Jurkat cell line lysates (35ug/lane). APG4A (arrow) was detected using the purified Pab.



APG4A Antibody (I378) (Cat. #AP1808c) western blot analysis in mouse kidney tissue lysates (35ug/lane). This demonstrates the APG4A antibody detected the APG4A protein (arrow).



Formalin-fixed and paraffin-embedded human cancer tissue reacted with the primary antibody, which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated. BC = breast carcinoma; HC = hepatocarcinoma.



Formalin-fixed and paraffin-embedded human skeletal muscle tissue reacted with Autophagy APG4A Antibody (C-term) (Cat.#AP1808c), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.

### **ATG4A Antibody - Background**

Macroautophagy is the major inducible pathway for the general turnover of cytoplasmic constituents in eukaryotic cells, it is also responsible for the degradation of active cytoplasmic enzymes and organelles during nutrient starvation. Macroautophagy involves the formation of double-membrane bound autophagosomes which enclose the cytoplasmic constituent targeted for degradation in a membrane bound structure, which then fuse with the lysosome (or vacuole) releasing a single-membrane bound autophagic bodies which are then degraded within the lysosome (or vacuole).

APG4A is a cysteine protease required for autophagy, which cleaves the C-terminal part of either MAP1LC3, GABARAPL2 or GABARAP, allowing the liberation of form I. A subpopulation of form I is subsequently converted to a smaller form (form II). Form II, with a revealed C-terminal glycine, is considered to be the phosphatidylethanolamine (PE)-conjugated form, and has the capacity for the binding to autophagosomes. Preferred substrate is GABARAPL2 followed by MAP1LC3A and GABARAP.

### **ATG4A Antibody - References**

- Baehrecke EH. Nat Rev Mol Cell Biol. 6(6):505-10. (2005)
- Lum JJ, et al. Nat Rev Mol Cell Biol. 6(6):439-48. (2005)
- Greenberg JT. Dev Cell. 8(6):799-801. (2005)
- Levine B. Cell. 120(2):159-62. (2005)
- Shintani T and Klionsky DJ. Science. 306(5698):990-5. (2004)