

**TRIM24 Antibody (N-term)**  
**Affinity Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP18114a**

**Specification**

---

**TRIM24 Antibody (N-term) - Product Information**

Application	WB,E
Primary Accession	<a href="#">O15164</a>
Other Accession	<a href="#">NP_003843.3</a>
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	116831
Antigen Region	16-45

**TRIM24 Antibody (N-term) - Additional Information**

**Gene ID** 8805

**Other Names**

Transcription intermediary factor 1-alpha, TIF1-alpha, 632-, E3 ubiquitin-protein ligase TRIM24, RING finger protein 82, Tripartite motif-containing protein 24, TRIM24, RNF82, TIF1, TIF1A

**Target/Specificity**

This TRIM24 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 16-45 amino acids from the N-terminal region of human TRIM24.

**Dilution**

WB~~1:1000

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

TRIM24 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

**TRIM24 Antibody (N-term) - Protein Information**

**Name** TRIM24

**Synonyms** RNF82, TIF1, TIF1A

**Function** Transcriptional coactivator that interacts with numerous nuclear receptors and coactivators and modulates the transcription of target genes. Interacts with chromatin depending on histone H3 modifications, having the highest affinity for histone H3 that is both unmodified at 'Lys-4' (H3K4me0) and acetylated at 'Lys-23' (H3K23ac). Has E3 protein-ubiquitin ligase activity. During the DNA damage response, participates in an autoregulatory feedback loop with TP53. Early in response to DNA damage, ATM kinase phosphorylates TRIM24 leading to its ubiquitination and degradation. After sufficient DNA repair has occurred, TP53 activates TRIM24 transcription, ultimately leading to TRIM24-mediated TP53 ubiquitination and degradation (PubMed:[24820418](#)). Plays a role in the regulation of cell proliferation and apoptosis, at least in part via its effects on p53/TP53 levels. Up- regulates ligand-dependent transcription activation by AR, GCR/NR3C1, thyroid hormone receptor (TR) and ESR1. Modulates transcription activation by retinoic acid (RA) receptors, including RARA. Plays a role in regulating retinoic acid-dependent proliferation of hepatocytes (By similarity). Participates also in innate immunity by mediating the specific 'Lys-63'-linked ubiquitination of TRAF3 leading to activation of downstream signal transduction of the type I IFN pathway (PubMed:[32324863](#)). Additionally, negatively regulates NLRP3/CASP1/IL-1beta-mediated pyroptosis and cell migration probably by ubiquitinating NLRP3 (PubMed:[33724611](#)).

#### Cellular Location


Nucleus. Cytoplasm. Mitochondrion. Note=Colocalizes with sites of active transcription. Predominantly nuclear. Translocated from nucleus to mitochondria to mediate antiviral immunity (PubMed:[32324863](#)). Localizes to sites of DNA damage (PubMed:[25593309](#)).

#### TRIM24 Antibody (N-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### TRIM24 Antibody (N-term) - Images



TRIM24 Antibody ( N-term) (Cat. #AP18114a) western blot analysis in U-937,mouse NIH/3T3 cell line lysates (35ug/lane).This demonstrates the TRIM24 antibody detected the TRIM24 protein (arrow).

#### **TRIM24 Antibody (N-term) - Background**

The protein encoded by this gene mediates transcriptional control by interaction with the activation function 2 (AF2) region of several nuclear receptors, including the estrogen, retinoic acid, and vitamin D3 receptors. The protein localizes to nuclear bodies and is thought to associate with chromatin and heterochromatin-associated factors. The protein is a member of the tripartite motif (TRIM) family. The TRIM motif includes three zinc-binding domains - a RING, a B-box type 1 and a B-box type 2 - and a coiled-coil region. Two alternatively spliced transcript variants encoding different isoforms have been described for this gene.

#### **TRIM24 Antibody (N-term) - References**

Kikuchi, M., et al. Biochim. Biophys. Acta 1793(12):1828-1836(2009)  
Lowe, J.K., et al. PLoS Genet. 5 (2), E1000365 (2009) :  
Sugiyama, N., et al. Mol. Cell Proteomics 6(6):1103-1109(2007)  
Matsuoka, S., et al. Science 316(5828):1160-1166(2007)  
Yondola, M.A., et al. J. Virol. 81(8):4264-4271(2007)