

**SIRT4 Antibody (N-term)**  
**Affinity Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP18208a**

**Specification**

---

**SIRT4 Antibody (N-term) - Product Information**

Application	WB,E
Primary Accession	<a href="#">O9Y6E7</a>
Other Accession	<a href="#">O1JQC6</a> , <a href="#">NP_036372.1</a>
Reactivity	Human, Mouse
Predicted	Bovine
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	35188
Antigen Region	67-95

**SIRT4 Antibody (N-term) - Additional Information**

**Gene ID** 23409

**Other Names**

NAD-dependent protein deacetylase sirtuin-4 {ECO:0000255|HAMAP-Rule:MF\_03161}, 351-  
{ECO:0000255|HAMAP-Rule:MF\_03161}, NAD-dependent ADP-ribosyltransferase sirtuin-4  
{ECO:0000255|HAMAP-Rule:MF\_03161}, 242- {ECO:0000255|HAMAP-Rule:MF\_03161}, Regulatory  
protein SIR2 homolog 4 {ECO:0000255|HAMAP-Rule:MF\_03161}, SIR2-like protein 4  
{ECO:0000255|HAMAP-Rule:MF\_03161}, SIRT4 {ECO:0000255|HAMAP-Rule:MF\_03161}, SIR2L4

**Target/Specificity**

This SIRT4 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 67-95 amino acids from the N-terminal region of human SIRT4.

**Dilution**

WB~~1:1000

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

SIRT4 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

**SIRT4 Antibody (N-term) - Protein Information**

**Name** SIRT4 {ECO:0000255|HAMAP-Rule:MF\_03161, ECO:0000312|HGNC:HGNC:14932}

**Function** Acts as a NAD-dependent protein lipoamidase, biotinylase, deacetylase and ADP-ribosyl transferase (PubMed:[16959573](#), PubMed:[17715127](#), PubMed:[24052263](#), PubMed:[25525879](#)). Catalyzes more efficiently removal of lipoyl- and biotinyl- than acetyl-lysine modifications (PubMed:[24052263](#), PubMed:[25525879](#)). Inhibits the pyruvate dehydrogenase complex (PDH) activity via the enzymatic hydrolysis of the lipoamide cofactor from the E2 component, DLAT, in a phosphorylation-independent manner (PubMed:[25525879](#)). Catalyzes the transfer of ADP-ribosyl groups onto target proteins, including mitochondrial GLUD1, inhibiting GLUD1 enzyme activity (PubMed:[16959573](#), PubMed:[17715127](#)). Acts as a negative regulator of mitochondrial glutamine metabolism by mediating mono ADP-ribosylation of GLUD1: expressed in response to DNA damage and negatively regulates anaplerosis by inhibiting GLUD1, leading to block metabolism of glutamine into tricarboxylic acid cycle and promoting cell cycle arrest (PubMed:[16959573](#), PubMed:[17715127](#)). In response to mTORC1 signal, SIRT4 expression is repressed, promoting anaplerosis and cell proliferation (PubMed:[23663782](#)). Acts as a tumor suppressor (PubMed:[23562301](#), PubMed:[23663782](#)). Also acts as a NAD-dependent protein deacetylase: mediates deacetylation of 'Lys-471' of MLYCD, inhibiting its activity, thereby acting as a regulator of lipid homeostasis (By similarity). Does not seem to deacetylate PC (PubMed:[23438705](#)). Controls fatty acid oxidation by inhibiting PPARA transcriptional activation (PubMed:[24043310](#)). Impairs SIRT1-PPARA interaction probably through the regulation of NAD(+) levels (PubMed:[24043310](#)). Down-regulates insulin secretion (PubMed:[17715127](#)).

#### Cellular Location

Mitochondrion matrix {ECO:0000255|HAMAP- Rule:MF\_03161, ECO:0000269|PubMed:16079181, ECO:0000269|PubMed:16959573, ECO:0000269|PubMed:17715127}

#### Tissue Location


Detected in vascular smooth muscle and striated muscle. Detected in insulin-producing beta-cells in pancreas islets of Langerhans (at protein level). Widely expressed. Weakly expressed in leukocytes and fetal thymus.


### SIRT4 Antibody (N-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### SIRT4 Antibody (N-term) - Images





SIRT4 Antibody (N-term) (Cat. #AP18208a) western blot analysis in mouse testis tissue lysates (35ug/lane). This demonstrates the SIRT4 antibody detected the SIRT4 protein (arrow).

### **SIRT4 Antibody (N-term) - Background**

This gene encodes a member of the sirtuin family of proteins, homologs to the yeast Sir2 protein. Members of the sirtuin family are characterized by a sirtuin core domain and grouped into four classes. The functions of human sirtuins have not yet been determined; however, yeast sirtuin proteins are known to regulate epigenetic gene silencing and suppress recombination of rDNA. Studies suggest that the human sirtuins may function as intracellular regulatory proteins with mono-ADP-ribosyltransferase activity. The protein encoded by this gene is included in class IV of the sirtuin family.

### **SIRT4 Antibody (N-term) - References**

Reiling, E., et al. *Eur. J. Hum. Genet.* 17(8):1056-1062(2009)  
Ahuja, N., et al. *J. Biol. Chem.* 282(46):33583-33592(2007)  
Yamamoto, H., et al. *Mol. Endocrinol.* 21(8):1745-1755(2007)  
Haigis, M.C., et al. *Cell* 126(5):941-954(2006)  
Scherer, S.E., et al. *Nature* 440(7082):346-351(2006)