

## **PCGF6 Antibody (Center)**

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP18850c

### **Specification**

# PCGF6 Antibody (Center) - Product Information

Application WB,E
Primary Accession Q9BYE7

Other Accession <u>Q5XI70</u>, <u>Q99NA9</u>, <u>NP\_001011663.1</u>

Reactivity
Predicted
Host
Clonality
Isotype
Calculated MW
Antigen Region

Human, Rat
Mouse
Rabbit
Polyclonal
Rabbit IgG
39047
190-217

# PCGF6 Antibody (Center) - Additional Information

#### **Gene ID 84108**

### **Other Names**

Polycomb group RING finger protein 6, Mel18 and Bmi1-like RING finger, RING finger protein 134, PCGF6, MBLR, RNF134

### Target/Specificity

This PCGF6 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 190-217 amino acids from the Central region of human PCGF6.

#### **Dilution**

WB~~1:1000

#### **Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

#### Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

#### **Precautions**

PCGF6 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

# PCGF6 Antibody (Center) - Protein Information

#### Name PCGF6



# Synonyms MBLR, RNF134

**Function** Transcriptional repressor (PubMed:12167161). May modulate the levels of histone H3K4Me3 by activating KDM5D histone demethylase (PubMed:17320162). Component of a Polycomb group (PcG) multiprotein PRC1-like complex, a complex class required to maintain the transcriptionally repressive state of many genes, including Hox genes, throughout development. PcG PRC1 complex acts via chromatin remodeling and modification of histones; it mediates monoubiquitination of histone H2A 'Lys-119', rendering chromatin heritably changed in its expressibility (PubMed:12167161). Within the PRC1-like complex, regulates RNF2 ubiquitin ligase activity (PubMed:26151332).

**Cellular Location** Nucleus


Tissue Location Ubiquitous.

### **PCGF6 Antibody (Center) - Protocols**

Provided below are standard protocols that you may find useful for product applications.


- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- <u>Immunofluorescence</u>
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

# PCGF6 Antibody (Center) - Images



PCGF6 Antibody (Center) (Cat.# AP18850c) western blot analysis in NCI-H292,rat liver,stomach and testis cell line lysates (35ug/lane). This demonstrates the PCGF6 antibody detected the PCGF6 protein (arrow).





All lanes: Anti-PCGF6 Antibody (Center) at 1:1000 dilution Lane 1: U-251 MG whole cell lysate Lane 2: K562 whole cell lysate Lane 3: NCI-H292 whole cell lysate Lysates/proteins at 20  $\mu$ g per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size: 39 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

## PCGF6 Antibody (Center) - Background

The protein encoded by this gene contains a RING finger motif, which is most closely related to those of polycomb group (PcG) proteins RNF110/MEL-18 and BMI1. PcG proteins are known to form protein complexes and function as transcription repressors. This protein has been shown to interact with some PcG proteins and act as a transcription repressor. The activity of this protein is found to be regulated by cell cycle dependent phosphorylation. Alternatively spliced transcript variants encoding different isoforms have been identified.

# **PCGF6 Antibody (Center) - References**

Lee, J.Y., et al. Cancer Res. 68(11):4201-4209(2008)
Zhang, J., et al. J. Biol. Chem. 283(12):7464-7469(2008)
Lee, M.G., et al. Cell 128(5):877-887(2007)
Zhong, N., et al. Biochem. Biophys. Res. Commun. 338(2):855-861(2005)
Deloukas, P., et al. Nature 429(6990):375-381(2004)