

Mouse Cdk6 Antibody (C-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP19011B

Specification

Mouse Cdk6 Antibody (C-term) - Product Information

Application	WB,E
Primary Accession	O64261
Other Accession	NP_034003.1
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	37028
Antigen Region	285-312

Mouse Cdk6 Antibody (C-term) - Additional Information

Gene ID 12571

Other Names

Cyclin-dependent kinase 6, CR2 protein kinase, CRK2, Cell division protein kinase 6, Serine/threonine-protein kinase PLSTIRE, Cdk6, Cdkn6, Crk2

Target/Specificity

This Mouse Cdk6 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 285-312 amino acids from the C-terminal region of mouse Cdk6.

Dilution

WB~~1:1000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Mouse Cdk6 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Mouse Cdk6 Antibody (C-term) - Protein Information

Name Cdk6

Synonyms Cdkn6, Crk2

Function Serine/threonine-protein kinase involved in the control of the cell cycle and differentiation; promotes G1/S transition. Phosphorylates pRB/RB1 and NPM1. Interacts with D-type G1 cyclins during interphase at G1 to form a pRB/RB1 kinase and controls the entrance into the cell cycle. Involved in initiation and maintenance of cell cycle exit during cell differentiation; prevents cell proliferation and negatively regulates cell differentiation, but is required for the proliferation of specific cell types (e.g. erythroid and hematopoietic cells). Essential for cell proliferation within the dentate gyrus of the hippocampus and the subventricular zone of the lateral ventricles. Required during thymocyte development. Promotes the production of newborn neurons, probably by modulating G1 length. Promotes, at least in astrocytes, changes in patterns of gene expression, changes in the actin cytoskeleton including loss of stress fibers, and enhanced motility during cell differentiation. Prevents myeloid differentiation by interfering with RUNX1 and reducing its transcription transactivation activity, but promotes proliferation of normal myeloid progenitors. Delays senescence. Promotes the proliferation of beta-cells in pancreatic islets of Langerhans (By similarity). May play a role in the centrosome organization during the cell cycle phases.

Cellular Location

Cytoplasm {ECO:0000250|UniProtKB:Q00534}. Nucleus {ECO:0000250|UniProtKB:Q00534}. Cell projection, ruffle {ECO:0000250|UniProtKB:Q00534}. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome {ECO:0000250|UniProtKB:Q00534} Note=Localized to the ruffling edge of spreading fibroblasts. Kinase activity only in nucleus (By similarity). Present in the cytosol and in the nucleus in interphase cells and at the centrosome during mitosis from prophase to telophase (By similarity). Localized to the cytosol of neurons and showed prominent staining around either side of the nucleus. {ECO:0000250|UniProtKB:Q00534, ECO:0000269|PubMed:23918663}

Tissue Location

Expressed in subgranular zone (SGZ) of the hippocampal dentate gyrus (DG) and the subventricular zone (SVZ) of the lateral ventricles whose neural precursor cells (NPC) give rise to dentate granule neurons and olfactory bulb (OB) interneurons, respectively. Expressed in the neuroepithelium of the cerebral cortex of the developing brain.


Mouse Cdk6 Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Mouse Cdk6 Antibody (C-term) - Images





Mouse Cdk6 Antibody (C-term) (Cat. #AP19011b) western blot analysis in HepG2 cell line lysates (35ug/lane). This demonstrates the Cdk6 antibody detected the Cdk6 protein (arrow).

Mouse Cdk6 Antibody (C-term) - Background

Cdk6 probably involved in the control of the cell cycle. Interacts with D-type G1 cyclins (By similarity).

Mouse Cdk6 Antibody (C-term) - References

- Puyol, M., et al. *Cancer Cell* 18(1):63-73(2010)
- Wiedemeyer, W.R., et al. *Proc. Natl. Acad. Sci. U.S.A.* 107(25):11501-11506(2010)
- Rivadeneira, D.B., et al. *Gastroenterology* 138(5):1920-1930(2010)
- Michaud, K., et al. *Cancer Res.* 70(8):3228-3238(2010)
- Choe, K.S., et al. *J. Biol. Chem.* 285(5):3044-3052(2010)