

NCAPG Antibody (N-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP19113a

Specification

NCAPG Antibody (N-term) - Product Information

Application	WB,E
Primary Accession	O9BPX3
Other Accession	NP_071741.2
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	114334
Antigen Region	117-145

NCAPG Antibody (N-term) - Additional Information

Gene ID 64151

Other Names

Condensin complex subunit 3, Chromosome-associated protein G, Condensin subunit CAP-G, hCAP-G, Melanoma antigen NY-MEL-3, Non-SMC condensin I complex subunit G, XCAP-G homolog, NCAPG, CAPG, NYMEL3

Target/Specificity

This NCAPG antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 117-145 amino acids from the N-terminal region of human NCAPG.

Dilution

WB~~1:1000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

NCAPG Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

NCAPG Antibody (N-term) - Protein Information

Name NCAPG

Synonyms CAPG, NYMEL3

Function Regulatory subunit of the condensin complex, a complex required for conversion of interphase chromatin into mitotic-like condense chromosomes. The condensin complex probably introduces positive supercoils into relaxed DNA in the presence of type I topoisomerases and converts nicked DNA into positive knotted forms in the presence of type II topoisomerases.

Cellular Location

Nucleus. Cytoplasm. Chromosome. Note=In interphase cells, the majority of the condensin complex is found in the cytoplasm, while a minority of the complex is associated with chromatin. A subpopulation of the complex however remains associated with chromosome foci in interphase cells. During mitosis, most of the condensin complex is associated with the chromatin. At the onset of prophase, the regulatory subunits of the complex are phosphorylated by CDK1, leading to condensin's association with chromosome arms and to chromosome condensation. Dissociation from chromosomes is observed in late telophase


Tissue Location

Highly expressed in testis.

NCAPG Antibody (N-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

NCAPG Antibody (N-term) - Images

NCAPG Antibody (N-term) (Cat. #AP19113a) western blot analysis in CEM cell line lysates (35ug/lane). This demonstrates the NCAPG antibody detected the NCAPG protein (arrow).

NCAPG Antibody (N-term) - Background

Regulatory subunit of the condensin complex, a complex required for conversion of interphase

chromatin into mitotic-like condense chromosomes. The condensin complex probably introduces positive supercoils into relaxed DNA in the presence of type I topoisomerases and converts nicked DNA into positive knotted forms in the presence of type II topoisomerases.

NCAPG Antibody (N-term) - References

Okada, Y., et al. Hum. Mol. Genet. 19(11):2303-2312(2010)
Zhao, J., et al. BMC Med. Genet. 11, 96 (2010) :
Murphy, L.A., et al. Biochem. Biophys. Res. Commun. 377(3):1007-1011(2008)
Gudbjartsson, D.F., et al. Nat. Genet. 40(5):609-615(2008)
Sugiyama, N., et al. Mol. Cell Proteomics 6(6):1103-1109(2007)

NCAPG Antibody (N-term) - Citations

- [Prognostic comparative genes predict targets for sorafenib combination therapies in hepatocellular carcinoma.](#)