

**APH1A Antibody (N-term)**  
**Affinity Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP19161A**

**Specification**

---

**APH1A Antibody (N-term) - Product Information**

Application	WB,E
Primary Accession	<a href="#">O96BI3</a>
Other Accession	<a href="#">O8BVF7</a> , <a href="#">NP_057106.2</a>
Reactivity	Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	28996
Antigen Region	41-70

**APH1A Antibody (N-term) - Additional Information**

**Gene ID** 51107

**Other Names**

Gamma-secretase subunit APH-1A, APH-1a, Aph-1alpha, Presenilin-stabilization factor, APH1A, PSF

**Target/Specificity**

This APH1A antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 41-70 amino acids from the N-terminal region of human APH1A.

**Dilution**

WB~~1:1000

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

APH1A Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

**APH1A Antibody (N-term) - Protein Information**

**Name** APH1A

**Synonyms** PSF

**Function** Non-catalytic subunit of the gamma-secretase complex, an endoprotease complex that catalyzes the intramembrane cleavage of integral membrane proteins such as Notch receptors and APP (amyloid- beta precursor protein) (PubMed:[12297508](#), PubMed:[12522139](#), PubMed:[12679784](#), PubMed:[12763021](#), PubMed:[25043039](#), PubMed:[26280335](#), PubMed:[30598546](#), PubMed:[30630874](#)). Required for normal gamma-secretase assembly (PubMed:[12471034](#), PubMed:[12522139](#), PubMed:[12763021](#), PubMed:[19369254](#)). The gamma-secretase complex plays a role in Notch and Wnt signaling cascades and regulation of downstream processes via its role in processing key regulatory proteins, and by regulating cytosolic CTNNB1 levels (Probable).

#### Cellular Location

Endoplasmic reticulum membrane; Multi-pass membrane protein. Golgi apparatus, Golgi stack membrane; Multi-pass membrane protein. Note=Predominantly located in the endoplasmic reticulum and in the cis-Golgi

#### Tissue Location


Widely expressed. Expressed in leukocytes, lung, placenta, small intestine, liver, kidney, spleen thymus, skeletal muscle, heart and brain. Isoform 1 and isoform 2 are nearly expressed at the same level.

### APH1A Antibody (N-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### APH1A Antibody (N-term) - Images



APH1A Antibody (N-term) (Cat. #AP19161a) western blot analysis in mouse liver tissue lysates (35ug/lane). This demonstrates the APH1A antibody detected the APH1A protein (arrow).

### APH1A Antibody (N-term) - Background

APH1 is a multipass transmembrane protein that interacts with presenilin (see PSEN1; MIM 104311) and nicastrin (APH2; MIM 605254) as a functional component of the gamma-secretase complex. The gamma-secretase complex is required for the intramembrane proteolysis of a number of membrane proteins, including the amyloid-beta precursor protein (APP; MIM 104760) and Notch (MIM 190198).

#### **APH1A Antibody (N-term) - References**

Rose, J.E., et al. Mol. Med. 16 (7-8), 247-253 (2010) :  
Mitsuishi, Y., et al. J. Biol. Chem. 285(20):14920-14931(2010)  
Chen, A.C., et al. J. Biol. Chem. 285(15):11378-11391(2010)  
Pardossi-Piquard, R., et al. J. Biol. Chem. 284(24):16298-16307(2009)  
Wang, Y., et al. Neurosci. Lett. 455(2):101-104(2009)