

**VAPA Antibody(C-term)**  
**Affinity Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP19461b****Specification**

---

**VAPA Antibody(C-term) - Product Information**

Application	WB,E
Primary Accession	<a href="#">O9P0L0</a>
Other Accession	<a href="#">O9Z270</a> , <a href="#">O9WV55</a> , <a href="#">NP_003565.4</a>
Reactivity	Human
Predicted	Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	27893
Antigen Region	182-210

**VAPA Antibody(C-term) - Additional Information****Gene ID** 9218**Other Names**

Vesicle-associated membrane protein-associated protein A, VAMP-A, VAMP-associated protein A, VAP-A, 33 kDa VAMP-associated protein, VAP-33, VAPA, VAP33

**Target/Specificity**

This VAPA antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 182-210 amino acids from the C-terminal region of human VAPA.

**Dilution**

WB~~1:1000

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

VAPA Antibody(C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

**VAPA Antibody(C-term) - Protein Information****Name** VAPA ([HGNC:12648](#))

## Synonyms VAP33

**Function** Endoplasmic reticulum (ER)-anchored protein that mediates the formation of contact sites between the ER and endosomes via interaction with FFAT motif-containing proteins such as STARD3 or WDR44 (PubMed:[32344433](#), PubMed:[33124732](#)). STARD3-VAPA interaction enables cholesterol transfer from the ER to endosomes (PubMed:[33124732](#)). Via interaction with WDR44 participates in neosynthesized protein export (PubMed:[32344433](#)). In addition, recruited to the plasma membrane through OSBPL3 binding (PubMed:[25447204](#)). The OSBPL3-VAPA complex stimulates RRAS signaling which in turn attenuates integrin beta-1 (ITGB1) activation at the cell surface (PubMed:[25447204](#)). With OSBPL3, may regulate ER morphology (PubMed:[16143324](#)). May play a role in vesicle trafficking (PubMed:[11511104](#), PubMed:[19289470](#)).

## Cellular Location

Endoplasmic reticulum membrane; Single-pass type IV membrane protein. Cell membrane; Single-pass type IV membrane protein. Cell junction, tight junction. Nucleus membrane {ECO:0000250|UniProtKB:Q9Z270}. Note=Present in the plasma membrane and in intracellular vesicles, together with SNARE proteins. May also associate with the cytoskeleton. Colocalizes with OCLN at the tight junction in polarized epithelial cells.

## Tissue Location


Ubiquitous.

## VAPA Antibody(C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

## VAPA Antibody(C-term) - Images



VAPA Antibody (C-term) (Cat. #AP19461b) western blot analysis in K562 cell line lysates (35ug/lane). This demonstrates the VAPA antibody detected the VAPA protein (arrow).

**VAPA Antibody(C-term) - Background**

The protein encoded by this gene is a type IV membrane protein. It is present in the plasma membrane and intracellular vesicles. It may also be associated with the cytoskeleton. This protein may function in vesicle trafficking, membrane fusion, protein complex assembly and cell motility. Alternative splicing occurs at this locus and two transcript variants encoding distinct isoforms have been identified.

**VAPA Antibody(C-term) - References**

Furuita, K., et al. J. Biol. Chem. 285(17):12961-12970(2010)  
Tuuf, J., et al. Biochem. Biophys. Res. Commun. 388(2):395-399(2009)  
Saita, S., et al. J. Biol. Chem. 284(20):13766-13777(2009)  
Prosser, D.C., et al. J. Cell. Sci. 121 (PT 18), 3052-3061 (2008) :  
Lohoff, F.W., et al. J Neural Transm 115(9):1339-1345(2008)