

CCR5 Antibody(Center)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP19617C

Specification

CCR5 Antibody(Center) - Product Information

Application	WB,E
Primary Accession	P51681
Other Accession	NP_000570.1
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	40524
Antigen Region	151-177

CCR5 Antibody(Center) - Additional Information

Gene ID 1234

Other Names

C-C chemokine receptor type 5, C-C CKR-5, CC-CKR-5, CCR-5, CCR5, CHEMR13, HIV-1 fusion coreceptor, CD195, CCR5, CMKBR5

Target/Specificity

This CCR5 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 151-177 amino acids from the Central region of human CCR5.

Dilution

WB~~1:1000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

CCR5 Antibody(Center) is for research use only and not for use in diagnostic or therapeutic procedures.

CCR5 Antibody(Center) - Protein Information

Name CCR5 ([HGNC:1606](#))

Synonyms CMKBR5

Function Receptor for a number of inflammatory CC-chemokines including CCL3/MIP-1-alpha, CCL4/MIP-1-beta and RANTES and subsequently transduces a signal by increasing the intracellular calcium ion level. May play a role in the control of granulocytic lineage proliferation or differentiation. Participates in T-lymphocyte migration to the infection site by acting as a chemotactic receptor (PubMed:[30713770](#)).

Cellular Location

Cell membrane; Multi-pass membrane protein

Tissue Location


Highly expressed in spleen, thymus, in the myeloid cell line THP-1, in the promyeloblastic cell line KG-1a and on CD4+ and CD8+ T-cells. Medium levels in peripheral blood leukocytes and in small intestine. Low levels in ovary and lung.

CCR5 Antibody(Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

CCR5 Antibody(Center) - Images



CCR5 Antibody (Center) (Cat. #AP19617c) western blot analysis in MDA-MB231 cell line lysates (35ug/lane). This demonstrates the CCR5 antibody detected the CCR5 protein (arrow).

CCR5 Antibody(Center) - Background

This gene encodes a member of the beta chemokine receptor family, which is predicted to be a seven transmembrane protein similar to G protein-coupled receptors. This protein is expressed by T cells and macrophages, and is known to be an important co-receptor for macrophage-tropic virus, including HIV, to enter host cells. Defective alleles of this gene have been associated

with the HIV infection resistance. The ligands of this receptor include monocyte chemoattractant protein 2 (MCP-2), macrophage inflammatory protein 1 alpha (MIP-1 alpha), macrophage inflammatory protein 1 beta (MIP-1 beta) and regulated on activation normal T expressed and secreted protein (RANTES). Expression of this gene was also detected in a promyeloblastic cell line, suggesting that this protein may play a role in granulocyte lineage proliferation and differentiation. This gene is located at the chemokine receptor gene cluster region. Two transcript variants encoding the same protein have been found for this gene.

CCR5 Antibody(Center) - References

Gerard, H.C., et al. Am. J. Med. Sci. 340(6):448-451(2010)
Hammad, M.M., et al. J. Biol. Chem. 285(45):34653-34664(2010)
Martens, H.A., et al. J. Rheumatol. 37(11):2226-2231(2010)
Weng, C.J., et al. Oral Oncol. 46(10):767-772(2010)
Mamtani, M., et al. PLoS ONE 5 (7), E11458 (2010) :