

EIF4EBP1 Antibody (T69)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP1981c**Specification**

EIF4EBP1 Antibody (T69) - Product Information

Application	WB, IHC-P,E
Primary Accession	Q13541
Other Accession	Q0P5A7 , NP_004086
Reactivity	Human
Predicted	Bovine
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	12580
Antigen Region	48-77

EIF4EBP1 Antibody (T69) - Additional Information**Gene ID** 1978**Other Names**

Eukaryotic translation initiation factor 4E-binding protein 1, 4E-BP1, eIF4E-binding protein 1, Phosphorylated heat- and acid-stable protein regulated by insulin 1, PHAS-I, EIF4EBP1

Target/Specificity

This EIF4EBP1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 48-77 amino acids from human EIF4EBP1.

DilutionWB~~1:1000
IHC-P~~1:10~50**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

EIF4EBP1 Antibody (T69) is for research use only and not for use in diagnostic or therapeutic procedures.

EIF4EBP1 Antibody (T69) - Protein Information**Name** EIF4EBP1

Function Repressor of translation initiation that regulates EIF4E activity by preventing its assembly into the eIF4F complex: hypophosphorylated form competes with EIF4G1/EIF4G3 and strongly binds to EIF4E, leading to repress translation. In contrast, hyperphosphorylated form dissociates from EIF4E, allowing interaction between EIF4G1/EIF4G3 and EIF4E, leading to initiation of translation. Mediates the regulation of protein translation by hormones, growth factors and other stimuli that signal through the MAP kinase and mTORC1 pathways.

Cellular Location


Cytoplasm. Nucleus. Note=Localization to the nucleus is unaffected by phosphorylation status. {ECO:0000250|UniProtKB:Q60876}

EIF4EBP1 Antibody (T69) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

EIF4EBP1 Antibody (T69) - Images



Western blot analysis of EIF4EBP1 (arrow) using rabbit polyclonal EIF4EBP1 Antibody (T69) (RB11407). 293 cell lysates (2 ug/lane) either nontransfected (Lane 1) or transiently transfected with the EIF4EBP1 gene (Lane 2) (Origene Technologies).



Formalin-fixed and paraffin-embedded human colon carcinoma tissue reacted with EIF4EBP1 Antibody (T69) (Cat.#AP1981c), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.

EIF4EBP1 Antibody (T69) - Background

EIF4EBP1 is one member of a family of translation repressor proteins. It directly interacts with eukaryotic translation initiation factor 4E (eIF4E), which is a limiting component of the multisubunit complex that recruits 40S ribosomal subunits to the 5' end of mRNAs. Interaction of this protein with eIF4E inhibits complex assembly and represses translation. This protein is phosphorylated in response to various signals including UV irradiation and insulin signaling, resulting in its dissociation from eIF4E and activation of mRNA translation.

EIF4EBP1 Antibody (T69) - References

Fonseca,B.D., J. Biol. Chem. 282 (34), 24514-24524 (2007)
Armengol,G., Cancer Res. 67 (16), 7551-7555 (2007)