

**RPS6 Antibody (N-term)**  
**Affinity Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP20175a**

**Specification**

---

**RPS6 Antibody (N-term) - Product Information**

Application	WB,E
Primary Accession	<a href="#">P62753</a>
Other Accession	<a href="#">P62755</a> , <a href="#">P62754</a> , <a href="#">O4R4K6</a> , <a href="#">Q5E995</a> , <a href="#">NP_001001.2</a> , <a href="#">G1TM55</a>
Reactivity	Human
Predicted	Bovine, Monkey, Mouse, Rabbit, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	28681
Antigen Region	1-30

**RPS6 Antibody (N-term) - Additional Information**

**Gene ID** 6194

**Other Names**

40S ribosomal protein S6, Phosphoprotein NP33, RPS6

**Target/Specificity**

This RPS6 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 1-30 amino acids from the N-terminal region of human RPS6.

**Dilution**

WB~~1:1000

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

RPS6 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

**RPS6 Antibody (N-term) - Protein Information**

**Name** RPS6 {ECO:0000303|PubMed:29563586, ECO:0000312|HGNC:HGNC:10429}

**Function** Component of the 40S small ribosomal subunit (PubMed:[23636399](#), PubMed:[8706699](#)). Plays an important role in controlling cell growth and proliferation through the selective translation of particular classes of mRNA (PubMed:[17220279](#)). Part of the small subunit (SSU) processome, first precursor of the small eukaryotic ribosomal subunit. During the assembly of the SSU processome in the nucleolus, many ribosome biogenesis factors, an RNA chaperone and ribosomal proteins associate with the nascent pre-rRNA and work in concert to generate RNA folding, modifications, rearrangements and cleavage as well as targeted degradation of pre-ribosomal RNA by the RNA exosome (PubMed:[34516797](#)).

#### Cellular Location


Cytoplasm. Nucleus, nucleolus

### RPS6 Antibody (N-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### RPS6 Antibody (N-term) - Images



RPS6 Antibody (N-term) (Cat. #AP20175a) western blot analysis in A431 cell line lysates (35ug/lane). This demonstrates the RPS6 antibody detected the RPS6 protein (arrow).

### RPS6 Antibody (N-term) - Background

Ribosomes, the organelles that catalyze protein synthesis, consist of a small 40S subunit and a large 60S subunit. Together these subunits are composed of 4 RNA species and approximately 80 structurally distinct proteins. This gene encodes a cytoplasmic ribosomal protein that is a component of the 40S subunit. The protein belongs to the S6E family of ribosomal proteins. It is the major substrate of protein kinases in the ribosome, with subsets of five C-terminal serine residues phosphorylated by different protein

kinases. Phosphorylation is induced by a wide range of stimuli, including growth factors, tumor-promoting agents, and mitogens. Dephosphorylation occurs at growth arrest. The protein may contribute to the control of cell growth and proliferation through the selective translation of particular classes of mRNA. As is typical for genes encoding ribosomal proteins, there are multiple processed pseudogenes of this gene dispersed through the genome.

#### **RPS6 Antibody (N-term) - References**

- Maggi, L.B. Jr., et al. Mol. Cell. Biol. 28(23):7050-7065(2008)  
Fujita, K., et al. Acta Neuropathol. 116(4):439-445(2008)  
Robledo, S., et al. RNA 14(9):1918-1929(2008)  
Glover, E.I., et al. Am. J. Physiol. Regul. Integr. Comp. Physiol. 295 (2), R604-R610 (2008) :  
Ma, X.M., et al. Cell 133(2):303-313(2008)