

**ETV7 Antibody (C-term)**  
**Affinity Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP20366b**

**Specification**

---

**ETV7 Antibody (C-term) - Product Information**

Application	WB,E
Primary Accession	<a href="#">O9Y603</a>
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	38998
Antigen Region	313-341

**ETV7 Antibody (C-term) - Additional Information**

**Gene ID** 51513

**Other Names**

Transcription factor ETV7, ETS translocation variant 7, ETS-related protein Tel2, Tel-related Ets factor, Transcription factor Tel-2, ETV7, TEL2, TELB, TREF

**Target/Specificity**

This ETV7 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 313-341 amino acids from the C-terminal region of human ETV7.

**Dilution**

WB~~1:2000

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

ETV7 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

**ETV7 Antibody (C-term) - Protein Information**

**Name** ETV7

**Synonyms** TEL2, TELB, TREF

**Function** Transcriptional repressor; binds to the DNA sequence 5'- CCGGAAGT-3'. Isoform A does not seem to have a repressor activity. Isoform C does not seem to have a repressor activity.

**Cellular Location**

Nucleus.

**Tissue Location**


Expressed in hematopoietic tissues.

**ETV7 Antibody (C-term) - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**ETV7 Antibody (C-term) - Images**



All lanes : Anti-ETV7 Antibody (C-term) at 1:2000 dilution Lane 1: A431 whole cell lysate Lane 2: Jurkat whole cell lysate Lane 3: K562 whole cell lysate Lane 4: NCI-H292 whole cell lysate Lane 5: U-87 MG whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 39 kDa Blocking/Dilution buffer: 5% NFD/MTBST.

**ETV7 Antibody (C-term) - Background**

Transcriptional repressor; binds to the DNA sequence 5'-CCGGAAGT-3'. Isoform A does not seem to have a repressor activity. Isoform C does not seem to have a repressor activity.