

CCNB2 Antibody (Center S92/T94)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP20412c

Specification

CCNB2 Antibody (Center S92/T94) - Product Information

Application	WB,E
Primary Accession	O95067
Other Accession	Q4R7A8
Reactivity	Human
Predicted	Monkey
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	45282
Antigen Region	72-99

CCNB2 Antibody (Center S92/T94) - Additional Information

Gene ID 9133

Other Names

G2/mitotic-specific cyclin-B2, CCNB2

Target/Specificity

This CCNB2 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 72-99 amino acids from the Central region of human CCNB2.

Dilution

WB~~1:1000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

CCNB2 Antibody (Center S92/T94) is for research use only and not for use in diagnostic or therapeutic procedures.

CCNB2 Antibody (Center S92/T94) - Protein Information

Name CCNB2


Function Essential for the control of the cell cycle at the G2/M (mitosis) transition.

CCNB2 Antibody (Center S92/T94) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

CCNB2 Antibody (Center S92/T94) - Images



All lanes : Anti-CCNB2 Antibody (Center S92/T94) at 1:1000 dilution Lane 1: HeLa whole cell lysate Lane 2: SW480 whole cell lysate Lysates/proteins at 20 μ g per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 45 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

CCNB2 Antibody (Center S92/T94) - Background

Essential for the control of the cell cycle at the G2/M (mitosis) transition.