

M Tfap2c Antibody (N-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP20531a

Specification

M Tfap2c Antibody (N-term) - Product Information

Application	WB,E
Primary Accession	O61312
Reactivity	Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	49137
Antigen Region	123-144

M Tfap2c Antibody (N-term) - Additional Information

Gene ID 21420

Other Names

Transcription factor AP-2 gamma, AP2-gamma, AP-22, Activating enhancer-binding protein 2 gamma, Tfap2c, Tcfap2c

Target/Specificity

This Mouse Tfap2c antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 123-144 amino acids from the N-terminal region of Mouse Tfap2c.

Dilution

WB~~1:1000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

M Tfap2c Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

M Tfap2c Antibody (N-term) - Protein Information

Name Tfap2c

Synonyms Tcfap2c

Function Sequence-specific DNA-binding protein that interacts with inducible viral and cellular enhancer elements to regulate transcription of selected genes. AP-2 factors bind to the consensus sequence 5'-GCCNNNGGC-3' and activate genes involved in a large spectrum of important biological functions including proper eye, face, body wall, limb and neural tube development. They also suppress a number of genes including MCAM/MUC18, C/EBP alpha and MYC.

Cellular Location

Nucleus.

Tissue Location


Expressed in lung, ovary and testis. Expressed in most squamous epithelia. Also, detected in several exocrine glands including the prostate, the preputial and salivary glands, serous glands of the tongue and ocular harderian glands

M Tfp2c Antibody (N-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

M Tfp2c Antibody (N-term) - Images



Mouse Tfp2c Antibody (N-term) (Cat. #AP20531a) western blot analysis in mouse heart tissue lysates (35ug/lane). This demonstrates the Mouse Tfp2c antibody detected the Mouse Tfp2c protein (arrow).

M Tfp2c Antibody (N-term) - Background

Sequence-specific DNA-binding protein that interacts with inducible viral and cellular enhancer elements to regulate transcription of selected genes. AP-2 factors bind to the consensus sequence

5'-GCCNNNGGC-3' and activate genes involved in a large spectrum of important biological functions including proper eye, face, body wall, limb and neural tube development. They also suppress a number of genes including MCAM/MUC18, C/EBP alpha and MYC.

M Tfp2c Antibody (N-term) - References

- Oulad-Abdelghani M., et al. Exp. Cell Res. 225:338-347(1996).
Carninci P., et al. Science 309:1559-1563(2005).
Church D.M., et al. PLoS Biol. 7:E1000112-E1000112(2009).
Mural R.J., et al. Submitted (JUL-2005) to the EMBL/GenBank/DDBJ databases.
Yahata T., et al. Genomics 80:601-613(2002).