

(DANRE) gfap Antibody (N-term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP20618a

Specification

(DANRE) gfap Antibody (N-term) - Product Information

Application	WB, IHC-P,E
Primary Accession	Q58EE9
Reactivity	Zebrafish
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	51249
Antigen Region	2-36

(DANRE) gfap Antibody (N-term) - Additional Information

Gene ID 30646

Other Names

Glial fibrillary acidic protein, GFAP, gfap

Target/Specificity

This (DANRE) gfap antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 2-36amino acids from the N-terminal region of Zebrafish(DANRE) gfap.

Dilution

WB~~1:1000

IHC-P~~1:25

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

(DANRE) gfap Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

(DANRE) gfap Antibody (N-term) - Protein Information

Name gfap

Cellular Location


Cytoplasm, cytoskeleton. Note=Associated with intermediate filaments

(DANRE) gfap Antibody (N-term) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

(DANRE) gfap Antibody (N-term) - Images



Western blot analysis of lysates from zebrafish brain, Zebrafish tissue lysate (from left to right), using (DANRE) gfap Antibody (N-term)(Cat. #AP20618a). AP20618a was diluted at 1:1000 at each lane. A goat anti-rabbit IgG H&L(HRP) at 1:5000 dilution was used as the secondary antibody. Lysates at 35ug per lane.



Immunohistochemical analysis of paraffin-embedded Z.muscle section using (DANRE) gfap Antibody (N-term)(Cat#AP20618a). AP20618a was diluted at 1:25 dilution. A peroxidase-conjugated goat anti-rabbit IgG at 1:400 dilution was used as the secondary antibody, followed by DAB staining.

(DANRE) gfap Antibody (N-term) - References

- Nielsen A.L.,et al.Gene 310:123-132(2003).
Emelyanov A.,et al.Submitted (SEP-2003) to the EMBL/GenBank/DDBJ databases.
Clark S.,et al.Submitted (AUG-1999) to the EMBL/GenBank/DDBJ databases.