

**HIST1H2BM Antibody (N-term)**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP20679a**

**Specification**

---

**HIST1H2BM Antibody (N-term) - Product Information**

|                   |                        |
|-------------------|------------------------|
| Application       | WB,E                   |
| Primary Accession | <a href="#">Q99879</a> |
| Reactivity        | Human                  |
| Host              | Rabbit                 |
| Clonality         | Polyclonal             |
| Isotype           | Rabbit IgG             |
| Calculated MW     | 13989                  |
| Antigen Region    | 10-44                  |

**HIST1H2BM Antibody (N-term) - Additional Information**

**Gene ID** 8342

**Other Names**

Histone H2B type 1-M, Histone H2Be, H2B/e, HIST1H2BM, H2BFE

**Target/Specificity**

This HIST1H2BM antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 10-44 amino acids from the N-terminal region of human HIST1H2BM.

**Dilution**

WB~~1:1000

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

HIST1H2BM Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

**HIST1H2BM Antibody (N-term) - Protein Information**

**Name** H2BC14 ([HGNC:4750](#))

**Function** Core component of nucleosome. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and

chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications of histones, also called histone code, and nucleosome remodeling.

#### Cellular Location


Nucleus. Chromosome.

### HIST1H2BM Antibody (N-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### HIST1H2BM Antibody (N-term) - Images



Western blot analysis of lysates from K562, PC-3 cell line and human skeletal muscle tissue lysate (from left to right), using HIST1H2BM Antibody (N-term) (Cat. #AP20679a). AP20679a was diluted at 1:1000 at each lane. A goat anti-rabbit IgG H&L (HRP) at 1:5000 dilution was used as the secondary antibody. Lysates at 35ug per lane.

### HIST1H2BM Antibody (N-term) - Background

Core component of nucleosome. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications of histones, also called histone code, and nucleosome remodeling.

### HIST1H2BM Antibody (N-term) - References

Albig W., et al. Hum. Genet. 101:284-294(1997).

Marzluff W.F., et al. Genomics 80:487-498(2002).  
Mungall A.J., et al. Nature 425:805-811(2003).  
Lubec G., et al. Submitted (MAR-2007) to UniProtKB.  
Cheung W.L., et al. Cell 113:507-517(2003).