

PRDX3 Antibody (C-term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP20728c

Specification

PRDX3 Antibody (C-term) - Product Information

Application	WB,E
Primary Accession	P30048
Reactivity	Human, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	27693

PRDX3 Antibody (C-term) - Additional Information

Gene ID 10935

Other Names

Thioredoxin-dependent peroxide reductase, mitochondrial, Antioxidant protein 1, AOP-1, HBC189, Peroxiredoxin III, Prx-III, Peroxiredoxin-3, Protein MER5 homolog, PRDX3, AOP1

Target/Specificity

This PRDX3 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 229-262 amino acids from the C-terminal region of human PRDX3.

Dilution

WB~~1:2000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

PRDX3 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

PRDX3 Antibody (C-term) - Protein Information

Name PRDX3

Synonyms AOP1

Function Thiol-specific peroxidase that catalyzes the reduction of hydrogen peroxide and organic

hydroperoxides to water and alcohols, respectively. Plays a role in cell protection against oxidative stress by detoxifying peroxides (PubMed:[17707404](#), PubMed:[29438714](#), PubMed:[33889951](#), PubMed:[7733872](#)). Acts synergistically with MAP3K13 to regulate the activation of NF-kappa-B in the cytosol (PubMed:[12492477](#)). Required for the maintenance of physical strength (By similarity).

Cellular Location


Mitochondrion. Cytoplasm. Early endosome. Note=Localizes to early endosomes in a RPS6KC1-dependent manner.

PRDX3 Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

PRDX3 Antibody (C-term) - Images



All lanes : Anti-PRDX3 Antibody (C-term) at 1:2000 dilution Lane 1: HeLa whole cell lysate Lane 2: HepG2 whole cell lysate Lane 3: MOLT-4 whole cell lysate Lane 4: MCF-7 whole cell lysate Lane 5: R. kidney whole lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 28 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

PRDX3 Antibody (C-term) - Background

Involved in redox regulation of the cell. Protects radical-sensitive enzymes from oxidative damage by a radical- generating system. Acts synergistically with MAP3K13 to regulate the activation of NF-kappa-B in the cytosol.

PRDX3 Antibody (C-term) - References

Tsuji K., et al. *Biochem. J.* 307:377-381(1995).

Ota T., et al. *Nat. Genet.* 36:40-45(2004).

Ebert L., et al. Submitted (MAY-2004) to the EMBL/GenBank/DDBJ databases.

Kalnine N., et al. Submitted (OCT-2004) to the EMBL/GenBank/DDBJ databases.

Deloukas P., et al. *Nature* 429:375-381(2004).