

**RSF1 Antibody (C-term)**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP20734c**

**Specification**

---

**RSF1 Antibody (C-term) - Product Information**

Application	<b>WB,E</b>
Primary Accession	<a href="#">O96T23</a>
Reactivity	<b>Human</b>
Host	<b>Rabbit</b>
Clonality	<b>Polyclonal</b>
Isotype	<b>Rabbit IgG</b>
Calculated MW	<b>163821</b>

**RSF1 Antibody (C-term) - Additional Information**

**Gene ID** 51773

**Other Names**

Remodeling and spacing factor 1, Rsf-1, HBV pX-associated protein 8, Hepatitis B virus X-associated protein, p325 subunit of RSF chromatin-remodeling complex, RSF1, HBXAP, XAP8

**Target/Specificity**

This RSF1 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 1355-1389 amino acids from the C-terminal region of human RSF1.

**Dilution**

WB~~1:1000

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

RSF1 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

**RSF1 Antibody (C-term) - Protein Information**

**Name** RSF1

**Synonyms** HBXAP, XAP8

**Function** Regulatory subunit of the ATP-dependent RSF-1 and RSF-5 ISWI chromatin-remodeling

complexes, which form ordered nucleosome arrays on chromatin and facilitate access to DNA during DNA-templated processes such as DNA replication, transcription, and repair (PubMed:[12972596](#), PubMed:[28801535](#)). Binds to core histones together with SMARCA5, and is required for the assembly of regular nucleosome arrays by the RSF-5 ISWI chromatin-remodeling complex (PubMed:[12972596](#)). Directly stimulates the ATPase activity of SMARCA1 and SMARCA5 in the RSF-1 and RSF-5 ISWI chromatin-remodeling complexes, respectively (PubMed:[28801535](#)). The RSF-1 ISWI chromatin remodeling complex has a lower ATP hydrolysis rate than the RSF-5 ISWI chromatin-remodeling complex (PubMed:[28801535](#)). The complexes do not have the ability to slide mononucleosomes to the center of a DNA template (PubMed:[28801535](#)). Facilitates transcription of hepatitis B virus (HBV) genes by the pX transcription activator. In case of infection by HBV, together with pX, it represses TNF-alpha induced NF-kappa-B transcription activation. Represses transcription when artificially recruited to chromatin by fusion to a heterogeneous DNA binding domain (PubMed:[11788598](#), PubMed:[11944984](#)).

#### Cellular Location

Nucleus Note=Localization is diffuse during mitosis (PubMed:[12972596](#)). Co-localizes with SMARCA5 in the nucleus (PubMed:[12972596](#))

#### Tissue Location


Ubiquitously expressed. Highly expressed in the heart, skeletal muscle, kidney and placenta (PubMed:[12972596](#)) Expressed at low levels in the brain and colon (PubMed:[12972596](#))

#### RSF1 Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### RSF1 Antibody (C-term) - Images



Western blot analysis of lysate from HeLa cell line, using RSF1 Antibody (C-term)(Cat. #AP20734c). AP20734c was diluted at 1:1000. A goat anti-rabbit IgG H&L(HRP) at 1:5000 dilution was used as the secondary antibody. Lysate at 35ug.

#### **RSF1 Antibody (C-term) - Background**

Required for assembly of regular nucleosome arrays by the RSF chromatin-remodeling complex. Facilitates transcription of hepatitis B virus (HBV) genes by the pX transcription activator. In case of infection by HBV, together with pX, it represses TNF- alpha induced NF-kappa-B transcription activation. Represses transcription when artificially recruited to chromatin by fusion to a heterogeneous DNA binding domain.

#### **RSF1 Antibody (C-term) - References**

Shamay M.,et al.Genomics 79:523-529(2002).  
Shamay M.,et al.J. Biol. Chem. 277:9982-9988(2002).  
Taylor T.D.,et al.Nature 440:497-500(2006).  
Mao Y.M.,et al.Submitted (APR-1998) to the EMBL/GenBank/DDBJ databases.  
Ota T.,et al.Nat. Genet. 36:40-45(2004).