

**JAK1 Antibody (C-term)**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP20770c**

**Specification**

---

**JAK1 Antibody (C-term) - Product Information**

Application	<b>WB,E</b>
Primary Accession	<a href="#">P23458</a>
Other Accession	<a href="#">P52332</a>
Reactivity	<b>Human, Mouse</b>
Host	<b>Rabbit</b>
Clonality	<b>Polyclonal</b>
Isotype	<b>Rabbit IgG</b>
Calculated MW	<b>133277</b>

**JAK1 Antibody (C-term) - Additional Information**

**Gene ID** 3716

**Other Names**

Tyrosine-protein kinase JAK1, Janus kinase 1, JAK-1, JAK1, JAK1A, JAK1B

**Target/Specificity**

This JAK1 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 866-899 amino acids from the C-terminal region of human JAK1.

**Dilution**

WB~~1:1000

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

JAK1 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

**JAK1 Antibody (C-term) - Protein Information**

**Name** JAK1

**Synonyms** JAK1A, JAK1B

**Function** Tyrosine kinase of the non-receptor type, involved in the IFN-alpha/beta/gamma signal

pathway (PubMed:[16239216](#), PubMed:[28111307](#), PubMed:[32750333](#), PubMed:[7615558](#), PubMed:[8232552](#)). Kinase partner for the interleukin (IL)-2 receptor (PubMed:[11909529](#)) as well as interleukin (IL)-10 receptor (PubMed:[12133952](#)). Kinase partner for the type I interferon receptor IFNAR2 (PubMed:[16239216](#), PubMed:[28111307](#), PubMed:[32750333](#), PubMed:[7615558](#), PubMed:[8232552](#)). In response to interferon-binding to IFNAR1-IFNAR2 heterodimer, phosphorylates and activates its binding partner IFNAR2, creating docking sites for STAT proteins (PubMed:[7759950](#)). Directly phosphorylates STAT proteins but also activates STAT signaling through the transactivation of other JAK kinases associated with signaling receptors (PubMed:[16239216](#), PubMed:[32750333](#), PubMed:[8232552](#)).

#### Cellular Location

Endomembrane system; Peripheral membrane protein. Note=Wholly intracellular, possibly membrane associated

#### Tissue Location


Expressed at higher levels in primary colon tumors than in normal colon tissue. The expression level in metastatic colon tumors is comparable to the expression level in normal colon tissue

### JAK1 Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### JAK1 Antibody (C-term) - Images



Western blot analysis of lysates from HT-29 cell line and mouse spleen tissue lysate (from left to right), using JAK1 Antibody (C-term) (Cat. #AP20770c). AP20770c was diluted at 1:1000 at each lane. A goat anti-rabbit IgG H&L (HRP) at 1:5000 dilution was used as the secondary antibody. Lysates at 35 µg per lane.

**JAK1 Antibody (C-term) - Background**

Tyrosine kinase of the non-receptor type, involved in the IFN-alpha/beta/gamma signal pathway. Kinase partner for the interleukin (IL)-2 receptor.

**JAK1 Antibody (C-term) - References**

Wilks A.F., et al. Mol. Cell. Biol. 11:2057-2065(1991).  
Totoki Y., et al. Submitted (MAR-2005) to the EMBL/GenBank/DDBJ databases.  
Gregory S.G., et al. Nature 441:315-321(2006).  
Mural R.J., et al. Submitted (SEP-2005) to the EMBL/GenBank/DDBJ databases.  
Lee S.-T., et al. Oncogene 8:3403-3410(1993).

**JAK1 Antibody (C-term) - Citations**

- [Roflumilast reverses polymicrobial sepsis-induced liver damage by inhibiting inflammation in mice.](#)