

DBH Antibody (Center)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP20809c

Specification

DBH Antibody (Center) - Product Information

| | |
|-------------------|------------------------|
| Application | WB,E |
| Primary Accession | P09172 |
| Reactivity | Human |
| Host | Rabbit |
| Clonality | Polyclonal |
| Isotype | Rabbit IgG |
| Calculated MW | 69065 |

DBH Antibody (Center) - Additional Information

Gene ID 1621

Other Names

Dopamine beta-hydroxylase, Dopamine beta-monoxygenase, Soluble dopamine beta-hydroxylase, DBH

Target/Specificity

This DBH antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 199-232 amino acids from the Central region of human DBH.

Dilution

WB~~1:1000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

DBH Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

DBH Antibody (Center) - Protein Information

Name DBH

Function Catalyzes the hydroxylation of dopamine to noradrenaline (also known as norepinephrine), and is thus vital for regulation of these neurotransmitters.

Cellular Location


[Soluble dopamine beta-hydroxylase]: Cytoplasmic vesicle, secretory vesicle lumen Cytoplasmic vesicle, secretory vesicle, chromaffin granule lumen. Secreted

DBH Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

DBH Antibody (Center) - Images



Western blot analysis of lysate from SH-SY5Y cell line, using DBH Antibody (Center)(Cat. #AP20809c). AP20809c was diluted at 1:1000. A goat anti-rabbit IgG H&L(HRP) at 1:10000 dilution was used as the secondary antibody. Lysate at 35ug.

DBH Antibody (Center) - Background

Conversion of dopamine to noradrenaline.

DBH Antibody (Center) - References

- Humphray S.J., et al. Nature 429:369-374(2004).
Lamouroux A., et al. EMBO J. 6:3931-3937(1987).
Kobayashi K., et al. Nucleic Acids Res. 17:1089-1102(1989).
Li B., et al. Biochem. J. 313:57-64(1996).
Liu T., et al. J. Proteome Res. 4:2070-2080(2005).