

Mouse Ror2 Antibody (C-term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP20846c

Specification

Mouse Ror2 Antibody (C-term) - Product Information

| | |
|-------------------|------------------------|
| Application | WB, IHC-P,E |
| Primary Accession | O9Z138 |
| Other Accession | Q01974 |
| Reactivity | Human |
| Host | Rabbit |
| Clonality | Polyclonal |
| Isotype | Rabbit IgG |
| Calculated MW | 105005 |

Mouse Ror2 Antibody (C-term) - Additional Information

Other Names

Tyrosine-protein kinase transmembrane receptor ROR2, mROR2, Neurotrophic tyrosine kinase, receptor-related 2, Ror2

Target/Specificity

This Mouse Ror2 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 756-790 amino acids from the C-terminal region of human Mouse Ror2.

Dilution

WB~~1:1000

IHC-P~~1:25

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Mouse Ror2 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Mouse Ror2 Antibody (C-term) - Protein Information

Name Ror2

Function Tyrosine-protein kinase receptor which may be involved in the early formation of the chondrocytes. It seems to be required for cartilage and growth plate development (PubMed:[10700181](#)). Phosphorylates YWHAB, leading to induction of osteogenesis and bone

formation. In contrast, has also been shown to have very little tyrosine kinase activity in vitro. May act as a receptor for wnt ligand WNT5A which may result in the inhibition of WNT3A-mediated signaling (By similarity).

Cellular Location


Cell membrane; Single-pass type I membrane protein

Mouse Ror2 Antibody (C-term) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Mouse Ror2 Antibody (C-term) - Images



Western blot analysis of lysates from HeLa, SH-SY5Y cell line (from left to right), using Mouse Ror2 Antibody (C-term)(Cat. #AP20846c). AP20846c was diluted at 1:1000 at each lane. A goat anti-rabbit IgG H&L(HRP) at 1:10000 dilution was used as the secondary antibody. Lysates at 20ug per lane.



Immunohistochemical analysis of paraffin-embedded M. stomach section using Mouse Ror2 Antibody (C-term)(Cat#AP20846c). AP20846c was diluted at 1:25 dilution. A undiluted biotinylated goat polyvalent antibody was used as the secondary, followed by DAB staining.

Mouse Ror2 Antibody (C-term) - Background

Tyrosine-protein kinase receptor which may be involved in the early formation of the chondrocytes. It seems to be required for cartilage and growth plate development. Phosphorylates YWHAB, leading to induction of osteogenesis and bone formation (By similarity).

Mouse Ror2 Antibody (C-term) - References

- Oishi I.,et al.Genes Cells 4:41-56(1999).
- Church D.M.,et al.PLoS Biol. 7:E1000112-E1000112(2009).
- DeChiara T.M.,et al.Nat. Genet. 24:271-274(2000).
- van Wijk N.V.,et al.Biochem. Biophys. Res. Commun. 390:211-216(2009).