

**Mouse Hmga2 Antibody (C-term)**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP20905c**

**Specification**

---

**Mouse Hmga2 Antibody (C-term) - Product Information**

Application	IF, WB,E
Primary Accession	<a href="#">P52927</a>
Other Accession	<a href="#">P52926</a>
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	11819

**Mouse Hmga2 Antibody (C-term) - Additional Information**

**Gene ID** 15364

**Other Names**

High mobility group protein HMGI-C, High mobility group AT-hook protein 2, Hmga2, Hmgic

**Target/Specificity**

This Mouse Hmga2 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 72-105 amino acids from the C-terminal region of Mouse Hmga2.

**Dilution**

IF~~1:25

WB~~1:1000

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

Mouse Hmga2 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

**Mouse Hmga2 Antibody (C-term) - Protein Information**

**Name** Hmga2

**Synonyms** Hmgic

**Function** Functions as a transcriptional regulator. Functions in cell cycle regulation through CCNA2. Plays an important role in chromosome condensation during the meiotic G2/M transition of spermatocytes. Plays a role in postnatal myogenesis, is involved in satellite cell activation (PubMed:[27446912](#)). Positively regulates IGF2 expression through PLAG1 and in a PLAG1-independent manner (By similarity).

**Cellular Location**

Nucleus.

**Tissue Location**


Expressed in mitotic spermatogonia, meiotic spermatocytes, and postmeiotic round spermatids (at protein level) (PubMed:14668482). Expressed in embryonic myogenic progenitor cells (PubMed:27446912).

**Mouse Hmga2 Antibody (C-term) - Protocols**


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**Mouse Hmga2 Antibody (C-term) - Images**



Fluorescent image of C2C12 cells stained with Mouse Hmga2 Antibody (C-term)(Cat#AP20905c). AP20905c was diluted at 1:25 dilution. An Alexa Fluor 488-conjugated goat anti-rabbit IgG at 1:400 dilution was used as the secondary antibody (green). Cytoplasmic actin was counterstained with Alexa Fluor® 555 conjugated with Phalloidin (red).



Western blot analysis of lysates from HepG2, NCCIT, mouse NIH/3T3, mouse F9 cell line (from left to right), using Hmga2 Antibody (C-term)(Cat. #AP20905c). AP20905c was diluted at 1:1000 at each lane. A goat anti-rabbit IgG H&L(HRP) at 1:10000 dilution was used as the secondary antibody. Lysates at 20ug per lane.

#### **Mouse Hmga2 Antibody (C-term) - Background**

Functions as a transcriptional regulator. Functions in cell cycle regulation through CCNA2. Plays an important role in chromosome condensation during the meiotic G2/M transition of spermatocytes.

#### **Mouse Hmga2 Antibody (C-term) - References**

- Manfioletti G.,et al.Nucleic Acids Res. 19:6793-6797(1991).
- Manfioletti G.,et al.Gene 167:249-253(1995).
- Zhou X.,et al.Nucleic Acids Res. 24:4071-4077(1996).
- Carninci P.,et al.Science 309:1559-1563(2005).
- Di Agostino S.,et al.Mol. Biol. Cell 15:1224-1232(2004).