

**INSRR Antibody (Center)**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP20928a**

**Specification**

---

**INSRR Antibody (Center) - Product Information**

Application	WB,E
Primary Accession	<a href="#">P14616</a>
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	143720

**INSRR Antibody (Center) - Additional Information**

**Gene ID** 3645

**Other Names**

Insulin receptor-related protein, IRR, IR-related receptor, Insulin receptor-related protein alpha chain, Insulin receptor-related protein beta chain, INSRR, IRR

**Target/Specificity**

This INSRR antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 668-702 amino acids from the Central region of human INSRR.

**Dilution**

WB~~1:4000

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

INSRR Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

**INSRR Antibody (Center) - Protein Information**

**Name** INSRR

**Synonyms** IRR

**Function** Receptor with tyrosine-protein kinase activity. Functions as a pH sensing receptor which is activated by increased extracellular pH. Activates an intracellular signaling pathway that involves IRS1 and AKT1/PKB.

**Cellular Location**


Membrane; Single-pass type I membrane protein.

### INSRR Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### INSRR Antibody (Center) - Images



Anti-INSRR Antibody (Center) at 1:4000 dilution + MCF-7 whole cell lysates Lysates/proteins at 20  $\mu$ g per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution Predicted band size : 144 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

### INSRR Antibody (Center) - Background

Receptor with tyrosine-protein kinase activity. Functions as a pH sensing receptor which is activated by increased extracellular pH. Activates an intracellular signaling pathway that involves IRS1 and AKT1/PKB.

### INSRR Antibody (Center) - References

- Haenze J.,et al.Horm. Metab. Res. 31:77-79(1999).  
Gregory S.G.,et al.Nature 441:315-321(2006).  
Mural R.J.,et al.Submitted (SEP-2005) to the EMBL/GenBank/DDBJ databases.  
Shier P.,et al.J. Biol. Chem. 264:14605-14608(1989).  
Greenman C.,et al.Nature 446:153-158(2007).