

CHMP3 Antibody (N-term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP21033a

Specification

CHMP3 Antibody (N-term) - Product Information

Application	WB,E
Primary Accession	Q9Y3E7
Other Accession	Q8CGS4 , Q9CQ10 , Q4R574 , Q58CS7
Reactivity	Human, Mouse
Predicted	Bovine, Monkey, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	25073

CHMP3 Antibody (N-term) - Additional Information

Gene ID 100526767;51652

Other Names

Charged multivesicular body protein 3, Chromatin-modifying protein 3, Neuroendocrine differentiation factor, Vacuolar protein sorting-associated protein 24, hVps24, CHMP3, CGI149, NEDF, VPS24

Target/Specificity

This CHMP3 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 28-61 amino acids from the N-terminal region of human CHMP3.

Dilution

WB~~1:1000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

CHMP3 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

CHMP3 Antibody (N-term) - Protein Information

Name CHMP3

Synonyms CGI149, NEDF, VPS24

Function Probable core component of the endosomal sorting required for transport complex III (ESCRT-III) which is involved in multivesicular bodies (MVBs) formation and sorting of endosomal cargo proteins into MVBs. MVBs contain intraluminal vesicles (ILVs) that are generated by invagination and scission from the limiting membrane of the endosome and mostly are delivered to lysosomes enabling degradation of membrane proteins, such as stimulated growth factor receptors, lysosomal enzymes and lipids. The MVB pathway appears to require the sequential function of ESCRT-O, -I, -II and -III complexes. ESCRT-III proteins mostly dissociate from the invaginating membrane before the ILV is released. The ESCRT machinery also functions in topologically equivalent membrane fission events, such as the terminal stages of cytokinesis and the budding of enveloped viruses (HIV-1 and other lentiviruses). ESCRT-III proteins are believed to mediate the necessary vesicle extrusion and/or membrane fission activities, possibly in conjunction with the AAA ATPase VPS4. Selectively binds to phosphatidylinositol 3,5-bisphosphate PtdIns(3,5)P2 and PtdIns(3,4)P2 in preference to other phosphoinositides tested. Involved in late stages of cytokinesis. Plays a role in endosomal sorting/trafficking of EGF receptor. Isoform 2 prevents stress-mediated cell death and accumulation of reactive oxygen species when expressed in yeast cells.

Cellular Location

Cytoplasm, cytosol. Membrane; Lipid-anchor. Endosome. Late endosome membrane.
Note=Localizes to the midbody of dividing cells


Tissue Location


Widely expressed. Expressed in heart, brain, placenta, lung, liver, skeletal muscle, kidney and pancreas

CHMP3 Antibody (N-term) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

CHMP3 Antibody (N-term) - Images



Western blot analysis of lysates from mouse 3T3-L1 cell line and mouse lung tissue (from left to right), using CHMP3 Antibody (N-term)(Cat. #AP21033a). AP21033a was diluted at 1:1000 at each lane. A goat anti-rabbit IgG H&L(HRP) at 1:10000 dilution was used as the secondary antibody. Lysates at 20ug per lane.



Western blot analysis of lysates from human brain and human heart tissue lysate (from left to right), using CHMP3 Antibody (N-term)(Cat. #AP21033a). AP21033a was diluted at 1:1000 at each lane. A goat anti-rabbit IgG H&L(HRP) at 1:10000 dilution was used as the secondary antibody. Lysates at 20ug per lane.

CHMP3 Antibody (N-term) - Background

Probable core component of the endosomal sorting required for transport complex III (ESCRT-III) which is involved in multivesicular bodies (MVBs) formation and sorting of endosomal cargo proteins into MVBs. MVBs contain intraluminal vesicles (ILVs) that are generated by invagination and scission from the limiting membrane of the endosome and mostly are delivered to lysosomes enabling degradation of membrane proteins, such as stimulated growth factor receptors, lysosomal enzymes and lipids. The MVB pathway appears to require the sequential function of ESCRT-O, -I,-II and -III complexes. ESCRT-III proteins mostly dissociate from the invaginating membrane before the ILV is released. The ESCRT machinery also functions in topologically equivalent membrane fission events, such as the terminal stages of cytokinesis and the budding of enveloped viruses (HIV-1 and other lentiviruses). ESCRT-III proteins are believed to mediate the necessary vesicle extrusion

and/or membrane fission activities, possibly in conjunction with the AAA ATPase VPS4. Selectively binds to phosphatidylinositol 3,5-bisphosphate PtdIns(3,5)P₂ and PtdIns(3,4)P₂ in preference to other phosphoinositides tested. Involved in late stages of cytokinesis. Plays a role in endosomal sorting/trafficking of EGF receptor. Isoform 2 prevents stress-mediated cell death and accumulation of reactive oxygen species when expressed in yeast cells.

CHMP3 Antibody (N-term) - References

- Wilson E.M., et al. J. Clin. Endocrinol. Metab. 86:4504-4511(2001).
Yan Q., et al. Exp. Cell Res. 304:265-273(2005).
Kemmer D., et al. BMC Genomics 7:48-48(2006).
Khoury C.M., et al. Gene 391:233-241(2007).
Lai C.-H., et al. Genome Res. 10:703-713(2000).