

EHD2 Antibody (C-term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP21050a

Specification

EHD2 Antibody (C-term) - Product Information

Application	WB, IHC-P-Leica,E
Primary Accession	O9NZN4
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Antigen Region	415-449

EHD2 Antibody (C-term) - Additional Information

Gene ID 30846

Other Names

EH domain-containing protein 2, PAST homolog 2, EHD2, PAST2

Target/Specificity

This EHD2 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 415-449 amino acids from the C-terminal region of human EHD2.

Dilution

WB~~1:500-1:2000
IHC-P-Leica~~1:500

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

EHD2 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

EHD2 Antibody (C-term) - Protein Information

Name EHD2 ([HGNC:3243](#))

Function ATP- and membrane-binding protein that controls membrane reorganization/tubulation upon ATP hydrolysis (By similarity). Plays a role in membrane trafficking between the plasma membrane and endosomes (PubMed:[17233914](#)). Important for the internalization of GLUT4.

Required for fusion of myoblasts to skeletal muscle myotubes. Required for normal translocation of FER1L5 to the plasma membrane (By similarity). Regulates the equilibrium between cell surface-associated and cell surface-dissociated caveolae by constraining caveolae at the cell membrane (PubMed:[25588833](#)).

Cellular Location

Cell membrane; Peripheral membrane protein {ECO:0000250|UniProtKB:Q8BH64}; Cytoplasmic side {ECO:0000250|UniProtKB:Q8BH64}. Membrane, caveola; Peripheral membrane protein {ECO:0000250|UniProtKB:Q8BH64}; Cytoplasmic side {ECO:0000250|UniProtKB:Q8BH64}. Endosome membrane {ECO:0000250|UniProtKB:Q4V8H8}; Peripheral membrane protein {ECO:0000250|UniProtKB:Q4V8H8}; Cytoplasmic side {ECO:0000250|UniProtKB:Q4V8H8}. Cytoplasm, cytosol {ECO:0000250|UniProtKB:Q8BH64}. Note=Colocalizes with GLUT4 in intracellular tubulovesicular structures that are associated with cortical F-actin. Colocalizes with FER1L5 at plasma membrane in myoblasts and myotubes. {ECO:0000250|UniProtKB:Q8BH64}

Tissue Location


Highly expressed in heart and moderately expressed in placenta, lung, and skeletal muscle.

EHD2 Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

EHD2 Antibody (C-term) - Images




All lanes : Anti-EHD2 Antibody (C-term) at 1:1000 dilution Lane 1: Human heart tissue lysate Lane 2: A549 whole cell lysate Lane 3: Human placenta tissue lysate Lane 4: HeLa whole cell lysate Lane 5: Mouse kidney tissue lysate Lane 6: Rat lung tissue lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted


band size : 61 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



All lanes : Anti-EHD2 Antibody (C-term) at 1:500-1:2000 dilution Lane 1: A549 whole cell lysate Lane 2: Human heart tissue lysate Lane 3: Mouse lung tissue lysate Lane 4: Rat lung tissue lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 61 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



Immunohistochemical analysis of paraffin-embedded human heart tissue using AP21050A performed on the Leica® BOND RXm. Tissue was fixed with formaldehyde at room temperature, antigen retrieval was by heat mediation with a EDTA buffer (pH9. 0). Samples were incubated with primary antibody(1:500) for 1 hours at room temperature. A undiluted biotinylated CRF Anti-Polyvalent HRP Polymer antibody was used as the secondary antibody.



Immunohistochemical analysis of paraffin-embedded human placenta tissue using AP21050A performed on the Leica® BOND RXm. Tissue was fixed with formaldehyde at room temperature, antigen retrieval was by heat mediation with a EDTA buffer (pH9. 0). Samples were incubated with primary antibody(1:500) for 1 hours at room temperature. A undiluted biotinylated CRF Anti-Polyvalent HRP Polymer antibody was used as the secondary antibody.

EHD2 Antibody (C-term) - Background

Plays a role in membrane reorganization in response to nucleotide hydrolysis. Binds to liposomes and deforms them into tubules. Plays a role in membrane trafficking between the plasma membrane and endosomes. Important for the internalization of GLUT4. Required for normal fusion of myoblasts to skeletal muscle myotubes. Required for translocation of FER1L5 to the plasma membrane. Binds ATP; does not bind GTP (By similarity).

EHD2 Antibody (C-term) - References

- Pohl U.,et al.Genomics 63:255-262(2000).
- Benjamin S.,et al.Submitted (DEC-2001) to the EMBL/GenBank/DDBJ databases.
- Ota T.,et al.Nat. Genet. 36:40-45(2004).
- Mural R.J.,et al.Submitted (JUL-2005) to the EMBL/GenBank/DDBJ databases.
- Aboulaich N.,et al.Biochem. J. 383:237-248(2004).