

**TMEM160 Antibody (N-term)**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP21058a**

**Specification**

---

**TMEM160 Antibody (N-term) - Product Information**

Application	WB,E
Primary Accession	<a href="#">O9NX00</a>
Other Accession	<a href="#">O9D938</a> , <a href="#">O24JY6</a>
Reactivity	Human, Mouse
Predicted	Bovine
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	19658
Antigen Region	32-64

**TMEM160 Antibody (N-term) - Additional Information**

**Gene ID** 54958

**Other Names**

Transmembrane protein 160, TMEM160

**Target/Specificity**

This TMEM160 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 32-64 amino acids from the N-terminal region of human TMEM160.

**Dilution**

WB~~1:1000

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

TMEM160 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

**TMEM160 Antibody (N-term) - Protein Information**

**Name** TMEM160 ([HGNC:26042](#))

**Cellular Location**


Mitochondrion inner membrane; Multi-pass membrane protein

### TMEM160 Antibody (N-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### TMEM160 Antibody (N-term) - Images



Western blot analysis of lysates from HT-29 cell line, mouse brain tissue lysate (from left to right), using TMEM160 Antibody (N-term)(Cat. #AP21058a). AP21058a was diluted at 1:1000 at each lane. A goat anti-rabbit IgG H&L(HRP) at 1:10000 dilution was used as the secondary antibody. Lysates at 20ug per lane.

### TMEM160 Antibody (N-term) - References

Ota T.,et al.Nat. Genet. 36:40-45(2004).  
Burkard T.R.,et al.BMC Syst. Biol. 5:17-17(2011).