

NR4A2 Antibody (Center)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP21092a**Specification**

NR4A2 Antibody (Center) - Product Information

Application	WB,E
Primary Accession	P43354
Other Accession	Q07917 , Q06219 , Q08E53
Reactivity	Human
Predicted	Bovine, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	66591

NR4A2 Antibody (Center) - Additional Information**Gene ID** 4929**Other Names**

Nuclear receptor subfamily 4 group A member 2, Immediate-early response protein NOT, Orphan nuclear receptor NURR1, Transcriptionally-inducible nuclear receptor, NR4A2, NOT, NURR1, TINUR

Target/Specificity

This NR4A2 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 380-412 amino acids from the Central region of human NR4A2.

Dilution

WB~~1:1000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

NR4A2 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

NR4A2 Antibody (Center) - Protein Information**Name** NR4A2**Synonyms** NOT, NURR1, TINUR

Function Transcriptional regulator which is important for the differentiation and maintenance of meso-diencephalic dopaminergic (mdDA) neurons during development (PubMed:[15716272](#), PubMed:[17184956](#)). It is crucial for expression of a set of genes such as SLC6A3, SLC18A2, TH and DRD2 which are essential for development of mdDA neurons (By similarity).

Cellular Location

Cytoplasm. Nucleus. Note=Mostly nuclear; oxidative stress promotes cytoplasmic localization

Tissue Location


Expressed in a number of cell lines of T-cell, B- cell and fibroblast origin. Strong expression in brain tissue

NR4A2 Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

NR4A2 Antibody (Center) - Images



Western blot analysis of lysate from human brain tissue lysate, using NR4A2 Antibody (Center)(Cat. #AP21092a). AP21092a was diluted at 1:1000. A goat anti-rabbit IgG H&L(HRP) at 1:10000 dilution was used as the secondary antibody. Lysate at 20ug.

NR4A2 Antibody (Center) - Background

Transcriptional regulator which is important for the differentiation and maintenance of meso-diencephalic dopaminergic (mdDA) neurons during development. It is crucial for expression of a set of genes such as SLC6A3, SLC18A2, TH and DRD2 which are essential for development of mdDA neurons (By similarity).

NR4A2 Antibody (Center) - References

Mages H.W.,et al.Mol. Endocrinol. 8:1583-1591(1994).
Ichinose H.,et al.Gene 230:233-239(1999).
Torii T.,et al.Gene 230:225-232(1999).
Hillier L.W.,et al.Nature 434:724-731(2005).
Mural R.J.,et al.Submitted (SEP-2005) to the EMBL/GenBank/DDBJ databases.