

KDM6B Antibody (Center)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP21160a

Specification

KDM6B Antibody (Center) - Product Information

Application	WB,E
Primary Accession	O15054
Reactivity	Mouse
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG
Calculated MW	176632

KDM6B Antibody (Center) - Additional Information

Gene ID 23135

Other Names

Lysine-specific demethylase 6B, 11411-, JmjC domain-containing protein 3, Jumonji domain-containing protein 3, Lysine demethylase 6B, KDM6B, JMJD3, KIAA0346

Target/Specificity

This KDM6B antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 879-912 amino acids from the Central region of human KDM6B.

Dilution

WB~~1:2000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

KDM6B Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

KDM6B Antibody (Center) - Protein Information

Name KDM6B

Synonyms JMJD3, KIAA0346

Function Histone demethylase that specifically demethylates 'Lys-27' of histone H3, thereby

playing a central role in histone code (PubMed:[17713478](#), PubMed:[17825402](#), PubMed:[17851529](#), PubMed:[18003914](#)). Demethylates trimethylated and dimethylated H3 'Lys-27' (PubMed:[17713478](#), PubMed:[17825402](#), PubMed:[17851529](#), PubMed:[18003914](#)). Plays a central role in regulation of posterior development, by regulating HOX gene expression (PubMed:[17851529](#)). Involved in inflammatory response by participating in macrophage differentiation in case of inflammation by regulating gene expression and macrophage differentiation (PubMed:[17825402](#)). Plays a demethylase-independent role in chromatin remodeling to regulate T-box family member-dependent gene expression by acting as a link between T-box factors and the SMARCA4- containing SWI/SNF remodeling complex (By similarity).

Cellular Location


Nucleus.

KDM6B Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

KDM6B Antibody (Center) - Images



Anti-KDM6B Antibody (Center) at 1:2000 dilution + mouse lung lysates Lysates/proteins at 20 μ g per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution Predicted band size : 177 kDa Blocking/Dilution buffer: 5% NFDN/TBST.

KDM6B Antibody (Center) - Background

Histone demethylase that specifically demethylates 'Lys- 27' of histone H3, thereby playing a central role in histone code. Demethylates trimethylated and dimethylated H3 'Lys-27'. Plays a central role in regulation of posterior development, by regulating HOX gene expression. Involved in

inflammatory response by participating in macrophage differentiation in case of inflammation by regulating gene expression and macrophage differentiation.

KDM6B Antibody (Center) - References

- Nagase T.,et al.DNA Res. 4:141-150(1997).
Nakajima D.,et al.DNA Res. 9:99-106(2002).
Zody M.C.,et al.Nature 440:1045-1049(2006).
Hu L.Y.,et al.Gene Expr. 13:179-189(2006).
De Santa F.,et al.Cell 130:1083-1094(2007).