

NPY2R Antibody (N-term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP21198a

Specification

NPY2R Antibody (N-term) - Product Information

Application	WB, IHC, FC,E
Primary Accession	P49146
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG
Calculated MW	42731
Antigen Region	29-65

NPY2R Antibody (N-term) - Additional Information

Gene ID 4887

Other Names

Neuropeptide Y receptor type 2, NPY2-R, NPY-Y2 receptor, Y2 receptor, NPY2R

Target/Specificity

This NPY2R antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 29-65 amino acids from the N-terminal region of human NPY2R.

Dilution

WB~~1:1000

IHC~~1:25

FC~~1:25

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

NPY2R Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

NPY2R Antibody (N-term) - Protein Information

Name NPY2R

Function Receptor for neuropeptide Y and peptide YY. The rank order of affinity of this receptor

for pancreatic polypeptides is PYY > NPY > PYY (3-36) > NPY (2-36) > [Ile-31, Gln-34] PP > [Leu-31, Pro-34] NPY > PP, [Pro-34] PYY and NPY free acid.

Cellular Location

Cell membrane; Multi-pass membrane protein.

Tissue Location


High levels in amygdala, corpus callosum, hippocampus and subthalamic nucleus. Also detectable in caudate nucleus, hypothalamus and substantia nigra

NPY2R Antibody (N-term) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


NPY2R Antibody (N-term) - Images




All lanes : Anti-NPY2R Antibody (N-term) at 1:2000 dilution Lane 1: mouse brain lysates Lane 2: rat brain lysates Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution Predicted band size : 43 kDa Blocking/Dilution buffer: 5% NFD/MTBST.



Anti-NPY2R Antibody (N-term) at 1:1000 dilution + human kidney lysates Lysates/proteins at 20 μ g per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution Predicted band size : 43 kDa Blocking/Dilution buffer: 5% NFD/MTBST.



AP21198a staining NPY2R in Human brain tissue sections by Immunohistochemistry (IHC-P - paraformaldehyde-fixed, paraffin-embedded sections). Tissue was fixed with formaldehyde and blocked with 3% BSA for 0.5 hour at room temperature; antigen retrieval was by heat mediation with a citrate buffer (pH6). Samples were incubated with primary antibody (1/25) for 1 hours at 37°C. A undiluted biotinylated goat polyvalent antibody was used as the secondary antibody.



Overlay histogram showing SH-SY5Y cells stained with AP21198a (green line). The cells were fixed with 4% paraformaldehyde (10 min) and then permeabilized with 90% methanol for 10 min. The cells were then incubated in 2% bovine serum albumin to block non-specific protein-protein interactions followed by the antibody (, 1:25 dilution) for 60 min at 37°C. The secondary antibody used was Alexa Fluor® 488 goat anti-rabbit IgG (H+L) (1583138) at 1/400 dilution for 40 min at 37°C. Isotype control antibody (blue line) was rabbit IgG1 (1µg/1x10⁶ cells) used under the same conditions. Acquisition of >10, 000 events was performed.

NPY2R Antibody (N-term) - Background

Receptor for neuropeptide Y and peptide YY. The rank order of affinity of this receptor for pancreatic polypeptides is PYY > NPY > PYY (3-36) > NPY (2-36) > [Ile-31, Gln-34] PP > [Leu- 31, Pro-34] NPY > PP, [Pro-34] PYY and NPY free acid.

NPY2R Antibody (N-term) - References

Gerald C.,et al.J. Biol. Chem. 270:26758-26761(1995).
Gehlert D.R.,et al.Mol. Pharmacol. 49:224-228(1996).
Rose P.M.,et al.J. Biol. Chem. 270:22661-22664(1995).
Yan H.,et al.Proc. Natl. Acad. Sci. U.S.A. 93:4661-4665(1996).
Ammar D.A.,et al.Genomics 38:392-398(1996).