

**(Mouse) Sox17 Antibody (C-term)**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP21226b**

**Specification**

---

**(Mouse) Sox17 Antibody (C-term) - Product Information**

Application	WB,E
Primary Accession	<a href="#">O61473</a>
Reactivity	Human, Mouse
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG
Antigen Region	353-387

**(Mouse) Sox17 Antibody (C-term) - Additional Information**

**Gene ID** 20671

**Other Names**

Transcription factor SOX-17, Sox17, Sox-17

**Target/Specificity**

This mouse Sox17 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 353-387 amino acids from the C-terminal region of mouse Sox17.

**Dilution**

WB~~1:1000

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

(Mouse) Sox17 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

**(Mouse) Sox17 Antibody (C-term) - Protein Information**

**Name** Sox17

**Synonyms** Sox-17

**Function** Acts as a transcription regulator that binds target promoter DNA and bends the DNA (PubMed:[19328208](#), PubMed:[24153254](#), PubMed:[8636240](#)). Binds to the sequences 5'-AACAAAT-'3

or 5'-AACAAAG-3' (PubMed:[8636240](#)). Modulates transcriptional regulation via WNT3A. Inhibits Wnt signaling. Promotes degradation of activated CTNNB1. Plays a key role in the regulation of embryonic development (PubMed:[11973269](#), PubMed:[17655922](#), PubMed:[24153254](#)). Required for normal development of the definitive gut endoderm (PubMed:[11973269](#)). Required for normal looping of the embryonic heart tube. Plays an important role in embryonic and postnatal vascular development, including development of arteries (PubMed:[24153254](#)). Plays an important role in postnatal angiogenesis, where it is functionally redundant with SOX18 (PubMed:[16895970](#)). Required for the generation and maintenance of fetal hematopoietic stem cells, and for fetal hematopoiesis (PubMed:[17655922](#)). Probable transcriptional activator in the premeiotic germ cells.

#### Cellular Location

Nucleus {ECO:0000255|PROSITE-ProRule:PRU00267, ECO:0000269|PubMed:20802155, ECO:0000269|PubMed:24153254, ECO:0000269|PubMed:8636240}

#### Tissue Location


Detected in lung and testis (PubMed:8636240). Detected in endothelial cells around small and large arteries in newborns and adults, but is barely detectable in veins (at protein level) (PubMed:24153254).

### (Mouse) Sox17 Antibody (C-term) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### (Mouse) Sox17 Antibody (C-term) - Images



Anti-(Mouse) Sox17 Antibody (C-term) at 1:2000 dilution + OVCAR3 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 45 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



Anti-(Mouse) Sox17 Antibody (C-term) at 1:1000 dilution + Mouse heart whole lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 45 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

#### **(Mouse) Sox17 Antibody (C-term) - Background**

Acts as transcription regulator that binds target promoter DNA and bends the DNA. Binds to the sequences 5'- AACAAAT-'3 or 5'-AACAAAG-3'. Modulates transcriptional regulation via WNT3A. Inhibits Wnt signaling. Promotes degradation of activated CTNNB1. Plays a key role in the regulation of embryonic development. Required for normal looping of the embryonic heart tube. Required for normal development of the definitive gut endoderm. Probable transcriptional activator in the premeiotic germ cells. Isoform 2 (T-SOX17) shows no DNA-binding activity.

#### **(Mouse) Sox17 Antibody (C-term) - References**

- Kanai Y., et al. J. Cell Biol. 133:667-681(1996).
- Carninci P., et al. Science 309:1559-1563(2005).
- Layfield R., et al. Submitted (FEB-1994) to the EMBL/GenBank/DDBJ databases.
- Kanai-Azuma M., et al. Development 129:2367-2379(2002).
- Kim I., et al. Cell 130:470-483(2007).