

PCGF6 Antibody (Center)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP21285c

Specification

PCGF6 Antibody (Center) - Product Information

Application	WB,E
Primary Accession	O9BYE7
Reactivity	Rat
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG
Calculated MW	39047
Antigen Region	215-249

PCGF6 Antibody (Center) - Additional Information

Gene ID 84108

Other Names

Polycomb group RING finger protein 6, Mel18 and Bmi1-like RING finger, RING finger protein 134, PCGF6, MBLR, RNF134

Target/Specificity

This PCGF6 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 215-249 amino acids from the Central region of human PCGF6.

Dilution

WB~~1:2000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

PCGF6 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

PCGF6 Antibody (Center) - Protein Information

Name PCGF6

Synonyms MBLR, RNF134

Function Transcriptional repressor (PubMed:[12167161](#)). May modulate the levels of histone H3K4Me3 by activating KDM5D histone demethylase (PubMed:[17320162](#)). Component of a Polycomb group (PcG) multiprotein PRC1-like complex, a complex class required to maintain the transcriptionally repressive state of many genes, including Hox genes, throughout development. PcG PRC1 complex acts via chromatin remodeling and modification of histones; it mediates monoubiquitination of histone H2A 'Lys-119', rendering chromatin heritably changed in its expressibility (PubMed:[12167161](#)). Within the PRC1-like complex, regulates RNF2 ubiquitin ligase activity (PubMed:[26151332](#)).

Cellular Location

Nucleus

Tissue Location


Ubiquitous.

PCGF6 Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

PCGF6 Antibody (Center) - Images



Anti-PCGF6 Antibody (Center) at 1:2000 dilution + rat testis lysates Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution Predicted band size : 39 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

PCGF6 Antibody (Center) - Background

Transcriptional repressor. May modulate the levels of histone H3K4Me3 by activating KDM5D

histone demethylase. Component of a Polycomb group (PcG) multiprotein PRC1-like complex, a complex class required to maintain the transcriptionally repressive state of many genes, including Hox genes, throughout development. PcG PRC1 complex acts via chromatin remodeling and modification of histones; it mediates monoubiquitination of histone H2A 'Lys-119', rendering chromatin heritably changed in its expressibility.

PCGF6 Antibody (Center) - References

- Akasaka T., et al. *Genes Cells* 7:835-850(2002).
Ota T., et al. *Nat. Genet.* 36:40-45(2004).
Deloukas P., et al. *Nature* 429:375-381(2004).
Lee M.G., et al. *Cell* 128:877-887(2007).
Maertens G.N., et al. *PLoS ONE* 4:E6380-E6380(2009).