

DYNC1H1 Antibody (C-term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP21385b

Specification

DYNC1H1 Antibody (C-term) - Product Information

Application	WB, IHC-P,E
Primary Accession	Q14204
Reactivity	Human, Mouse
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG
Antigen Region	4202-4236

DYNC1H1 Antibody (C-term) - Additional Information

Gene ID 1778

Other Names

Cytoplasmic dynein 1 heavy chain 1, Cytoplasmic dynein heavy chain 1, Dynein heavy chain, cytosolic, DYNC1H1, DHC1, DNCH1, DNCL, DNECL, DYHC, KIAA0325

Target/Specificity

This DYNC1H1 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 4202-4236 amino acids from the C-terminal region of human DYNC1H1.

Dilution

WB~~1:2000

IHC-P~~1:25

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

DYNC1H1 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

DYNC1H1 Antibody (C-term) - Protein Information

Name DYNC1H1 ([HGNC:2961](#))

Function Cytoplasmic dynein 1 acts as a motor for the intracellular retrograde motility of vesicles and organelles along microtubules. Dynein has ATPase activity; the force-producing power stroke

is thought to occur on release of ADP. Plays a role in mitotic spindle assembly and metaphase plate congression (PubMed:[27462074](#)).

Cellular Location


Cytoplasm, cytoskeleton

DYNC1H1 Antibody (C-term) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


DYNC1H1 Antibody (C-term) - Images



All lanes : Anti-DYNC1H1 Antibody (C-term) at 1:2000 dilution Lane 1: U-87 MG whole cell lysates Lane 2: PC-3 whole cell lysates Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution Predicted band size : 532 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



All lanes : Anti-DYNC1H1 Antibody (C-term) at 1:2000 dilution Lane 1: A431 whole cell lysate Lane 2: PC-3 whole cell lysate Lane 3: mouse brain lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 532 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



AP21385b staining DYNC1H1 in human testis sections by Immunohistochemistry (IHC-P - paraformaldehyde-fixed, paraffin-embedded sections). Tissue was fixed with formaldehyde and blocked with 3% BSA for 0.5 hour at room temperature; antigen retrieval was by heat mediation with a citrate buffer (pH6). Samples were incubated with primary antibody (1/25) for 1 hours at 37°C. A undiluted biotinylated goat polyvalent antibody was used as the secondary antibody.

DYNC1H1 Antibody (C-term) - Background

Cytoplasmic dynein 1 acts as a motor for the intracellular retrograde motility of vesicles and organelles along microtubules. Dynein has ATPase activity; the force-producing power stroke is thought to occur on release of ADP.

DYNC1H1 Antibody (C-term) - References

Nagase T., et al. DNA Res. 4:141-150(1997).
Ohara O., et al. Submitted (AUG-2005) to the EMBL/GenBank/DDBJ databases.
Yamakawa H., et al. Submitted (JAN-2007) to the EMBL/GenBank/DDBJ databases.

Vaisberg E.A., et al. J. Cell Biol. 133:831-842(1996).
Vaisberg E.A., et al. J. Cell Biol. 123:849-858(1993).